

SUMMARY FISHING RECORD
Form 3600-63

Department of Natural Resources

County Oneida	Waters Ginty Lake MWBC: 1571300
Sampling Objective Bass & panfish assessment	Number and Locations of Stations (Habitat)
Period Fished (Dates)	Miles Actually Shocked = 2.0 Acres = 131 Total Miles of Shoreline = 2.8 Total Miles of Shockable Shoreline = 2.0
DATE 05/19/2011	Source GPS Topo Topo GPS

GEAR


Boomshocker (Hours) gamefish (entire shoreline) all species (half-mile reach)	1.0 0.3	Time X Night Day		
Visual Hours	Time of Day	Haul Seine (Length)	Mesh Size	Area Covered
Angling (Hours)	Time of Day	Mini fyke (No. of Net)	Mesh Size	Depth
Minnow Seine (No. of Hauls)	Area Covered	Gill Net (No. of Feet x of Lifts)	Mesh Size	Depth
Other (Hours or Lifts) Boomshocker(s): 1 Dip Netter(s): 2	Mini-boomshocker(s): Dip Netter(s):	Characteristics Water Temp: 59		

FISHING RESULTS

Species	No.	Modal Size(s)	Size Range	Catch/Unit
Walleye		-	-	
Largemouth Bass	2	-	13.0 - 13.9	1.94 / hour 1.00 / mile
Smallmouth Bass		-	-	
Muskellunge		-	-	
Northern Pike	11	-	15.0 - 32.2	10.65 /hour 5.50 / mile
Bluegill	95	6.75	2.5 - 7.9	335.29 /hour 190 / mile
Black Crappie	55	4.75, 8.25	2.5 - 8.9	194.12 /hour 110 / mile
Yellow Perch	21	2.75	2.0 - 8.9	74.12 /hour 42 / mile
Pumpkinseed	2	-	3.0 - 6.9	7.06 /hour 4 / mile
BluegillxPumpkinseed hybrids	2	-	6.0 - 6.9	7.06 /hour 4 / mile
Yellow Bullhead	1	-	10.5 - 10.9	3.53 /hour 2 / mile
Goldern Shiner	12	5.25	4.5 - 7.4	42.35 /hour 24 / mile
White Sucker	7	-	7.0 - 15.4	24.71 /hour 14 / mile

OBSERVATIONS

1) Mortality: None.	2) Weather: cloudy, warm.	3) Reliabilty: High
4) Stocking: None.	5) Adverse Conditions: None.	
6) Comments: collected all gamefish along the entire shoreline. All species along one 0.5-mile station.		
Signed (Compiler)	John Kubisiak	Date 11/15/11



**WISCONSIN DNR
FISHERIES INFORMATION SHEET**

LAKE: Ginty Lake

COUNTY: ONEIDA

YEAR: 2011

The Department of Natural Resources surveyed Ginty Lake Oneida County with a boomshocker on May 19, 2011. Ginty Lake is a drainage lake with dark-stained, soft water and has a surface area of 131 acres, 2.8 miles of shoreline and a maximum depth of 12 feet. The lake is divided into two basins by a gravel constriction which is spanned by a railroad bridge. Low clearance on the bridge kept us from electrofishing the 0.8 miles of shoreline in the south basin.




Figure 1. Length frequency distribution of 11 northern pike captured during a survey of Ginty Lake, Oneida County, May 19, 2011.

Black Crappie

Panfish and nongame fishes were only collected along two half-mile reaches of shoreline. We captured 55 black crappie, ranging from 2.5 to 8.9 inches. Peak abundance was at 8.25 inches.

Northern Pike

We captured 11 northern pike, including 6 fish over 25 inches. The largest pike was 32.2 inches in length.




Figure 2. Length frequency distribution of 55 black crappie captured during a survey of Ginty Lake, Oneida County, May 19, 2011.




Figure 3. Length frequency distribution of 95 bluegill captured during a survey of Ginty Lake, Oneida County, May 19, 2011.

Yellow perch

We captured 21 yellow perch, but most were small juveniles with only 3 fish larger than 7 inches.

Bluegill

We captured 95 bluegill, with very good numbers of fish up to 7.5 inches, but only one larger fish.




Figure 4. Length frequency distribution of 21 yellow perch captured during a survey of Ginty Lake, Oneida County, May 19, 2011.

Other Species

We captured 14 species of fish in our electroshocking sample of Ginty Lake. In addition to the species listed above, we caught 2 pumpkinseed, 2 bluegillxpumpkinseed hybrids, 1 yellow bullhead, 12 golden shiner and 7 white sucker.

John Kubisiak, Fisheries Biologist
Wisconsin Department of Natural Resources
107 Sutliff Avenue
Rhineland, WI 54501
(715) 365-8919
Email: JohnF1.Kubisiak@wisconsin.gov

MWBCODE: 1571300
 YEAR: 2011
 LAKE: Ginty Lake
 COUNTY: ONEIDA

Crew: Timler
 Kubisiak
 Nelson

M

Largemouth Bass		Northern Pike		Black Crappie		Yellow Perch		Bluegill	
Length Interval	Number								
<6				0.0		0.0		0.0	
7				0.5		0.5		0.5	
8		9		1.0		1.0		1.0	
9		10		1.5		1.5		1.5	
10		11		2.0		2.0	1	2.0	
11		12		2.5	1	2.5	9	2.5	2
12		13		3.0		3.0		3.0	3
13	2	14		3.5		3.5	1	3.5	1
14		15	2	4.0	1	4.0	1	4.0	5
15		16		4.5	8	4.5	1	4.5	8
16		17	1	5.0	4	5.0		5.0	4
17		18	1	5.5	1	5.5		5.5	9
18		19		6.0		6.0	2	6.0	21
19		20	1	6.5		6.5	3	6.5	25
20		21		7.0	3	7.0		7.0	16
21		22		7.5	14	7.5	2	7.5	1
22		23		8.0	17	8.0		8.0	
23		24		8.5	6	8.5	1	8.5	
24		25	1	9.0		9.0		9.0	
25+		26	1	9.5		9.5		9.5	
		27		10.0		10.0		10.0	
		28	1	10.5		10.5		10.5	
		29	2	11.0		11.0		11.0	
		30		11.5		11.5		11.5	
		31		12.0		12.0		12.0	
		32	1	12.5		12.5		12.5	
		33		13.0		13.0		13.0	
		34		13.5		13.5		13.5	
		35+		14.0		14.0		14.0	
				14.5		14.5		14.5	
				15.0		15.0		15.0	

Total	2	Total	11	Total	55	Total	21	Total	95
--------------	----------	--------------	-----------	--------------	-----------	--------------	-----------	--------------	-----------

Other species	count	lengths
Observed		
Golden Shiner	12	
White Sucker	7	
Pumpkinseed	2	
BluegillxPumpkinseed hybrids	2	
Yellow Bullhead	1	