ARCADIS

Table 1. Office Indoor Air Analytical Results, Madison-Kipp Corporation, 201 Waubesa Street, Madison, Wisconsin.

Sample Name Sample Date	Wisconsin Vapor Action Level ^{1,2} Non-Residential Indoor Air	Calculated Screening Level ³ Non-Residential Sub-Slab		AA-01		CS-01 12/13/2013	CS-02 12/13/2013	IA-01 12/13/2013
			12/13/2013	1/24/2014	7/18/2014			
VOCs								
cis-1,2-Dichloroethene	NE	NE	< 0.03	< 0.032	< 0.034	< 0.035	< 0.033	< 0.035
trans-1,2-Dichloroethene	65	6,500	< 0.15	<0.16	< 0.017	< 0.17	<0.17	< 0.17
Trichloroethene	1.6	160	< 0.03	< 0.032	< 0.034	< 0.035	< 0.033	< 0.035
Tetrachloroethene	27	2,700	< 0.03	< 0.032	< 0.034	0.24	0.11	< 0.035
Vinyl Chloride	11	1,100	< 0.015	< 0.016	< 0.017	< 0.017	< 0.017	< 0.017

All units presented in parts per billion by volume (ppbv).

Exceeds the Wisconsin Vapor Action Level for Non-Residential Indoor Air.

100 Exceeds the Calculated Screening Level for Non-Residential Sub-Slab Vapor.

- Constituent not detected above noted laboratory detection limit.
- Wisconsin Vapor Action Levels are the lower of the United States Environmental Protection Agency (U.S. EPA) Residential Screening Levels (RSLs) based on either a target cancer risk of 10-5 or a noncancer HQ of 1, and are presented in WDNR's *Indoor Air Vapor Action Levels for Various VOCS* (November 2013).
- Section VI A 1 of Addressing Vapor Intrusion at Remediation and Redevelopment Sites in Wisconsin (WDNR, 2010), accessed at: http://dnr.wi.gov/files/PDF/pubs/rr/RR800.pdf
- For large commercial/industrial buildings, the following attenuation factor was used: 0.01 for sub-slab vapor to indoor air [Section VI A 2 of WDNR (2010)].
- AA Ambient Air Sample.
- CS Crawlspace Sample.
- IA Indoor Air Sample.
- NE Not Established.
- VOC Volatile organic compound.

Table 1. Office Indoor Air Analytical Results, Madison-Kipp Corporation, 201 Waubesa Street, Madison, Wisconsin.

Sample Name	IA-02	IA-03	IA-04	IA-05		
Sample Date	12/13/2013	12/13/2013	12/13/2013	12/13/2013	1/24/2014	7/18/2014
VOCs						
cis-1,2-Dichloroethene	< 0.035	< 0.033	0.074	0.47	0.059	0.035
trans-1,2-Dichloroethene	<0.18	<0.17	<0.17	<0.16	<0.16	< 0.17
Trichloroethene	< 0.035	< 0.033	0.3	3	0.28	0.26
Tetrachloroethene	< 0.035	0.14	5.8	21	2.0	1.7
Vinyl Chloride	<0.018	<0.017	<0.017	<0.016	<0.016	0.024

All units presented in parts per billion by volume (ppbv).

Exceeds the Wisconsin Vapor Action Level for Non-Residential Indoor Air.

100 Exceeds the Calculated Screening Level for Non-Residential Sub-Slab Vapor.

- < Constituent not detected above noted laboratory detection limit.
- Wisconsin Vapor Action Levels are the lower of the United States Environmental Protection Agency (U.S. EPA) Residential Screening Levels (RSLs) based on either a target cancer risk of 10-5 or a noncancer HQ of 1, and are presented in WDNR's *Indoor Air Vapor Action Levels for Various VOCS* (November 2013).
- 2 Section VI A 1 of Addressing Vapor Intrusion at Remediation and Redevelopment Sites in Wisconsin (WDNR, 2010), accessed at: http://dnr.wi.gov/files/PDF/pubs/rr/RR800.pdf
- For large commercial/industrial buildings, the following attenuation factor was used: 0.01 for sub-slab vapor to indoor air [Section VI A 2 of WDNR (2010)].
- AA Ambient Air Sample.
- CS Crawlspace Sample.
- IA Indoor Air Sample.
- NE Not Established.
- VOC Volatile organic compound.