

LTA: 212Ka08NAME:
Oula Washed MoraineACRES:
34287.457SQUARE MILES:
53.574**DESCRIPTION:**

The characteristic landform pattern is undulating outwash plain and moraine. Soils are predominantly excessively drained loamy sand over outwash or acid loamy sand debris flow. Common habitat types include AQV-S, PMV-P, AC, and wetland.

CLIMATE

CODE	PERCENT
15	39
19	58
25	2

GEOLOGYBEDROCK TYPE DESCRIPTION
Igneous, Metamorphic, and Volcanic RockAVERAGE DEPTH
TO BEDROCK
10000BEDROCK DEPTH
DESCRIPTION
Bedrock is between 100 feet and 50 feet of the land surface**GEOMORPHOLOGY**GEOMORPHOLOGY PROCESS
Glacial Meltwater and Till DepositionTOPOGRAPHY
UndulatingSURFACE
Outwash Plain and Hummocky Moraine**SOIL INFORMATION**SOIL ASSOCIATIONS
Vilas-Keweenaw-SultzSOIL DESCRIPTION
Excessively drained and well drained sandy soils with a loamy sand surface over non-calcareous sand outwash, loamy sand till, or loamy lacustrine.SURFACE TEXTURES
LSGENERAL TEXTURES
SandyFAMILY TEXTURES
SDRAINAGE CLASSES
ED-WDPARENT MATERIAL
Outwash-Till-Lacustrine**KOTAR'S HABITAT**

HABITAT 1 HABITAT 2 HABITAT 3 HABITAT 4 HABITAT 5 HABITAT 6

*Listed in order of probability occurrence, with each having an occurrence of 10% or greater

WISCLAND LAND COVER

COVER TYPE CLASS	ACRES	PERCENT
Agricultural Land	32	0
Bare Land	174	1
Forested Wetland	5021	15
Grassland	2062	6
Nonforested Wetland	3322	10
Open Water	93	0
Shrubland	94	0
Upland Broad-leaved Deciduous	17039	50
Upland Coniferous Forest	697	2
Upland Mixed Deciduous/Conifer	5747	17

FINLEY'S PRESETTLEMENT VEGETATION INFORMATION

CODE	PERCENT	DEFINITION
BF	30	White Spruce, Balasam Fir, Tamarack, White Cedar, White Birch, Aspen
NH/P	7	Sugar Maple, Yellow Birch, White Pine, Red Pine
PW/PR	37	White Pine, Red Pine
ABW/P	4	Aspen, White Birch, Pine
SC	22	Swamp Conifers - White Cedar, Black Spruce, Tamarack, Hemlock

HYDROLOGY

PERENNIAL STREAMS	INTERMITTENT STREAMS	OPEN WATER	MARSH ACRES	DEPTH TO AQUIFER	SURFACE DRAINAGE
27 Miles	17 Miles	36 Acres	4251 Acres	20'-50'	deranged