

LTA: 212Qd06NAME:
Mead MarshACRES:
24924.236SQUARE MILES:
38.944**DESCRIPTION:**

The characteristic landform pattern is nearly level marsh. Soils are very poorly drained muck. Common habitat type is forested lowland.

CLIMATE

CODE	PERCENT
54	85
68	15

GEOLOGYBEDROCK TYPE DESCRIPTION
Igneous, Metamorphic, and Volcanic RockAVERAGE DEPTH
TO BEDROCK
5005BEDROCK DEPTH
DESCRIPTION
Bedrock is between 50 feet and 5 feet of the land surface**GEOMORPHOLOGY**GEOMORPHOLOGY PROCESS
Organic DepositionTOPOGRAPHY
Nearly LevelSURFACE
Swamp**SOIL INFORMATION**SOIL ASSOCIATIONS
Seelyeville-Markey-CathroSOIL DESCRIPTION
Very poorly drained nonacid muck, some over sand outwash or loamy alluvium.SURFACE TEXTURES
MKGENERAL TEXTURES
MuckyFAMILY TEXTURES
SP-S-LDRAINAGE CLASSES
VPDPARENT MATERIAL
Organic-Outwash-Alluvium**KOTAR'S HABITAT**

HABITAT 1	HABITAT 2	HABITAT 3	HABITAT 4	HABITAT 5	HABITAT 6
Lowland					

*Listed in order of probability occurrence, with each having an occurrence of 10% or greater

WISCLAND LAND COVER

COVER TYPE CLASS	ACRES	PERCENT
Agricultural Land	825	3
Bare Land	2	0
Forested Wetland	7497	30
Grassland	427	2
Nonforested Wetland	11027	44
Open Water	1837	7
Shrubland	5	0
Upland Broad-leaved Deciduous	3166	13
Upland Coniferous Forest	47	0
Upland Mixed Deciduous/Conifer	94	0

FINLEY'S PRESETTLEMENT VEGETATION INFORMATION

CODE	PERCENT	DEFINITION
Hydrographic	2	Water
HH/P	7	Hemlock, Sugar Maple, Yellow Birch, White Pine, Red Pine
NH/P	11	Sugar Maple, Yellow Birch, White Pine, Red Pine
OO	0	Oak - Black, White, and Burr Oak
SC	80	Swamp Conifers - White Cedar, Black Spruce, Tamarack, Hemlock

HYDROLOGY

PERENNIAL STREAMS	INTERMITTENT STREAMS	OPEN WATER	MARSH ACRES	DEPTH TO AQUIFER	SURFACE DRAINAGE
39 Miles	12 Miles	2820 Acres	13113 Acres	0'-20'	dendritic