

LTA: 212Xe13NAME:
Rhineland MorainesACRES:
37273.453SQUARE MILES:
58.24**DESCRIPTION:**

The characteristic landform pattern is undulating moraine and outwash plain. Soils are predominantly moderately well drained loam over acid loamy sand till or outwash. Common habitat types include ATM, TMC, forested lowland, and AViO.

CLIMATECODE PERCENT
33 100**GEOLOGY**BEDROCK TYPE DESCRIPTION
Igneous, Metamorphic, and Volcanic RockAVERAGE DEPTH
TO BEDROCK
10050BEDROCK DEPTH
DESCRIPTION
Bedrock is greater than 100 feet from the land surface**GEOMORPHOLOGY**

GEOMORPHOLOGY PROCESS

TOPOGRAPHY

SURFACE

SOIL INFORMATION

SOIL ASSOCIATIONS

SOIL DESCRIPTION

SURFACE TEXTURES

GENERAL TEXTURES

FAMILY TEXTURES

DRAINAGE CLASSES

PARENT MATERIAL

KOTAR'S HABITAT

HABITAT 1 HABITAT 2 HABITAT 3 HABITAT 4 HABITAT 5 HABITAT 6

*Listed in order of probability occurrence,
with each having an occurrence of 10% or
greater**WISCLAND LAND COVER**

COVER TYPE CLASS	ACRES	PERCENT
Agricultural Land	719	2
Bare Land	738	2
Forested Wetland	6883	20
Grassland	1065	3
High Intensity Urban Area	12	0
Low Intensity Urban Area	4	0
Nonforested Wetland	4056	12
Open Water	3664	11
Shrubland	789	2
Upland Broad-leaved Deciduous	11430	34
Upland Coniferous Forest	2671	8
Upland Mixed Deciduous/Conifer	1551	5

FINLEY'S PRESETTLEMENT VEGETATION INFORMATION

CODE	PERCENT	DEFINITION
Hydrographic	8	Water
HH/P	49	Hemlock, Sugar Maple, Yellow Birch, White Pine, Red Pine
NH/P	3	Sugar Maple, Yellow Birch, White Pine, Red Pine
PW/PR	8	White Pine, Red Pine
PJ/OX	1	Jack Pine, Scrub Oak and Barrens
ABW/P	1	Aspen, White Birch, Pine
SC	27	Swamp Conifers - White Cedar, Black Spruce, Tamarack, Hemlock

HYDROLOGY

PERENNIAL STREAMS	INTERMITTENT STREAMS	OPEN WATER	MARSH ACRES	DEPTH TO AQUIFER	SURFACE DRAINAGE
30 Miles	0 Miles	3663 Acres	10624 Acres	0'-20'	