

Aquatic Plant Management

Aquatic plants are a critical component in an aquatic ecosystem. Any management of an ecosystem can have negative or even detrimental effects on the whole ecosystem. Therefore, the practice of managing aquatic plants should not be taken lightly. The concept of Aquatic Plant Management (APM) is highly variable since different aquatic resource users want different things. Ideal management to one individual may mean providing prime fish habitat, for another it may be to remove surface vegetation for boating. The practice of APM is also highly variable. There are numerous APM strategies designed to achieve different plant management goals. Some are effective on a small scale, but ineffective in larger situations. Others can only be used for specific plants or during certain times of the growing season. Of course, the types of plants that are to be managed will also help determine which APM alternatives are feasible. The following paragraphs discuss the APM methods used today. The discussion is largely adopted from *Managing Lakes and Rivers, North American Lake Management Society, 2001*, supplemented with other applicable current resources and references. The methods summarized here are largely for management of rooted aquatic plants, not algae. While some methods may also have effects on nuisance algae blooms, the focus is submergent rooted aquatic macrophytes. This information is provided to allow the user to gain a basic understanding of the APM method, it is not designed to an all-inclusive APM decision-making matrix. APM alternatives can be divided into the following categories: Physical Controls, Chemical Controls, and Biological Controls.

Physical Controls

Physical APM controls include various methods to prevent growth or remove part or all of the aquatic plant. Both manual and mechanical techniques are employed. Physical APM methods include:

- ▲ Hand pulling
- ▲ Hand cutting
- ▲ Bottom barriers
- ▲ Light limitation (dyes, covers)
- ▲ Mechanical harvesting
- ▲ Hydorraking/rototilling
- ▲ Suction Dredging
- ▲ Dredging
- ▲ Drawdown

Each of these methods are described below. The costs, benefits, and drawbacks of each APM strategy are provided.

Hand Pulling: This method involves digging out the entire unwanted plant including stems and roots with a hand tool such as a spade. This method is highly selective and suitable for shallow areas for removing invasive species that have not become well established. This technique is obviously not for use on large dense beds of nuisance aquatic plants. It is best used in areas less than 3 feet, but can be used in deeper areas with divers using scuba and snorkeling equipment. It can also be used in combination with the suction dredge method. In Wisconsin, hand pulling may be completed outside a designated sensitive area without a permit but is limited to 30 feet of shoreline frontage. Removal of exotic species is not limited to 30 feet.

Advantages: This technique results in immediate clearing of the water column of nuisance plants. When a selective technique is desired in a shallow, small area, hand pulling is a good choice. It is also useful in sensitive areas where disruption must be minimized.

Disadvantages: This method is labor intensive. Disturbing the substrate may affect fish habitat, increase turbidity, and may promote phosphorus re-suspension and subsequent algae blooms.

Costs: The costs are highly variable. There is practically no cost using volunteers or lakeshore landowners to remove unwanted plants, however, using divers to remove plants can get relatively expensive. Hand pulling labor can range from \$400 to \$800 per acre.

Hand Cutting: This is another manual method where the plants are cut below the water surface. Generally the roots are not removed. Tools such as rakes, scythes or other specialized tools are pulled through the plant beds by boat or several people. This method is not as selective as hand pulling. This method is well suited for small areas near docks and piers. Plant material must be removed from the water. In Wisconsin, hand cutting may be completed outside a designated sensitive area without a permit but is limited to 30 feet of shoreline frontage. Removal of exotic species is not limited to 30 feet.

Advantages: This technique results in immediate clearing of the water column of nuisance plants. Costs are minimal.

Disadvantages: This is also a fairly time consuming and labor intensive option. Since the technique does not remove the entire plant (leaves root system and part of plant), it may not result in long-term reductions in growth. This technique is not species specific and results in all aquatic plants being removed from the water column.

Costs: The costs range from minimal for volunteers using hand equipment up to over \$1,000 for a hand-held mechanized cutting implement. Hand cutting labor can range from \$400 to \$800 per acre.

Bottom Barriers: A barrier material is applied over the lake bottom to prevent rooted aquatics from growing. Natural barriers such as clay, silt, and gravel can be used although eventually plants may root in these areas again. Artificial materials can also be used for bottom barriers and anchored to the substrate. Barrier materials include burlap, nylon, rubber, polyethylene, polypropylene, and fiberglass. Barriers include both solid and porous forms. A permit is required to place any fill or barrier structure on the substrate of a waterbody. This method is well suited for areas near docks, piers, and beaches. Periodic maintenance may be required to remove accumulated silt or rooting fragments from the barrier.

Advantages: This technique does not result in production of plant fragments. Properly installed, it can provide immediate and multiple year relief.

Disadvantages: This is a non-selective option, all plants beneath the barrier will be affected. Some materials are costly and installation is labor intensive. Other disadvantages include limited material durability, gas accumulation beneath the cover, or possible re-growth of plants from above or below the cover. Fish and invertebrate habitat is disrupted with this technique. Anchored barriers can be difficult to remove.

Costs: A 20 foot x 60 foot panel cost \$265, while a 30 foot x 50 foot panel cost \$375 (this does not include installation costs). Costs for materials vary from \$0.15 per square foot (ft²) to over \$0.35/ ft². The costs for installation range from \$0.25 to \$0.50/ ft². Barriers can cost \$20,000 to \$50,000 per acre.

Light Limitation: Limiting the available light in the water column can prevent photosynthesis and plant growth. Dark colored dyes and surface covers have been used to accomplish light limitation. Dyes are effective in shallow water bodies where their concentration can be kept at a desired concentration and loss through dilution is less. This method is well suited for small, shallow water bodies with no outlets such as private ponds.

Surface covers can be a useful tool in small areas such as docks and beaches. While they can interfere with aquatic recreation, they can be timed to produce results and not affect summer recreation uses.

Advantages: Dyes are non-toxic to humans and aquatic organisms. No special equipment is required for application. Light limitation with dyes or covers method may be selective to shade tolerant species. In addition to submerged macrophyte control, it can also control the algae growth.

Disadvantages: The application of water column dyes is limited to shallow water bodies with no outlets. Repeated dye treatments may be necessary. The dyes may not control peripheral or shallow-water rooted plants. This technique must be initiated before aquatic plants start to grow. Covers inhibit gas exchange with the atmosphere.

Costs: Costs for a commercial dye and application range from \$100 to \$500 per acre.

Mechanical Harvesting: Mechanical harvesters are essentially cutters mounted on barges that cut aquatic plants at a desired depth. Maximum cutting depths range from 5 to 8 feet with a cutting width of 6.5 to 12 feet. Cut plant materials require collection and removal from the water. Conventional harvesters combine cutting, collecting, storing, and transporting cut vegetation into one piece of equipment. Transport barges and shoreline conveyors are also available to remove the cut vegetation. The cut plants must be removed from the water body. The equipment needs are dictated by severity of the aquatic plant problem. Contract harvesting services are available in lieu of purchasing used or new equipment. Trained staff will be necessary to operate a mechanical harvester. To achieve maximum removal of plant material, harvesting is usually completed during the summer months while submergent vegetation is growing to the surface. The duration of control is variable and re-growth of aquatic plants is common. Factors such as timing of harvest, water depth, depth of cut, and timing can influence the effectiveness of a harvesting operation. Harvesting is suited for large open areas with dense stands of exotic or nuisance plant species. Permits are now required in Wisconsin to use a mechanical harvester.

Advantages: Harvesting provides immediate visible results. Harvesting allows plant removal on a larger scale than other options. Harvesting provides flexible area control. In other words, the harvester can be moved to where it is needed and used to target problem areas. This technique has the added benefit of removing the plant material from the water body and therefore also eliminates a possible source of nutrients often released during fall decay of aquatic plants. While removal of nutrients through plant harvesting has not been quantified, it can be important in aquatic ecosystem with low nutrient inputs.

Disadvantages: Drawbacks of harvesting include: limited depth of operation, not selective within the application area, and expensive equipment costs.

Harvesting also creates plant fragments, which can be a concern since certain plants have the ability to reproduce from a plant fragment (e.g. Eurasian watermilfoil). Plant fragments may re-root and spread a problem plant to other areas. Harvesting can have negative effects on non-target plants, young of year fish, and invertebrates. The harvesting will require trained operators and maintenance of equipment. Also, a disposal site or landspreading program will be needed for harvested plants.

Costs:

Costs for a harvesting operation are highly variable dependant on program scale. New harvesters range from \$40,000 for small machines to over \$100,000 for large, deluxe models. Costs vary considerably, depending on the model, size, and options chosen. Specially designed units are available, but may cost more. The equipment can last 10 to 15 years. A grant for ½ the equipment cost can be obtained from the Wisconsin Waterways Commission and a loan can be obtained for the remaining capital investment. Operation costs include insurance, fuel, spare parts, and payroll. Historical harvesting values have been reported at \$200 up to \$1,500 per acre. A survey of recent Wisconsin harvesting operations reported costs to be between \$100/acre and \$200/acre.

A used harvester can be purchased for \$10,000 to \$20,000. Maintenance costs are typically higher.

Contract harvesting costs approximately \$125/per hour plus mobilization to the water body. Contractors can typically harvest ¼ to ½ acre per hour for an estimated cost of \$250 to \$500/per acre.

Hydroraking/rototilling: Hydroraking is the use of a boat or barge mounted machine with a rake that is lowered to the bottom and dragged. The tines of the rake rip out roots of aquatic plants. Rototilling, or rotovation, also rips out root masses but uses a mechanical rotating head with tines instead of a rake. Harvesting may need to be completed in conjunction with these methods to gather floating plant fragments. This application would best be used where nuisance populations are well established and prevention of stem fragments is not critical. A permit would be required for this type of aquatic plant management and would only be issued in limited cases of extreme infestations of nuisance vegetation. In Wisconsin, this method is not looked upon favorably or at all by the WDNR.

Advantages:

These methods have the potential for significant reductions in aquatic plant growth. These methods can remove the plant stems and roots, resulting in thorough plant disruption. Hydroraking/rototilling can be completed in “off season” months avoiding interference with summer recreation activities.

Disadvantages:

Hydroraking/rototilling are not selective and may destroy substrate habitat important to fish and invertebrates. Suspension of sediments will increase turbidity and release nutrients trapped in bottom sediments into the water column potentially causing algal blooms. These methods can cause floating plant and root fragments, which may re-root and spread the problem. Hydroraking/rototilling are expensive and not likely to be permitted by regulatory agencies.

Costs: Bottom tillage costs vary according to equipment, treatment scale, and plant density. For soft vegetation costs can range from \$2,000 to \$4,000 per acre. For dense, rooted masses, costs can be up to \$10,000 per acre. Contract bottom tillage reportedly ranges from \$1,200 to \$1,700 per acre (Washington Department of Ecology, 1994).

Suction Dredging: Suction dredging uses a small boat or barge with portable dredges and suction heads. Scuba divers operate the suction dredge and can target removal of whole plants, seeds, and roots. This method may be applied in conjunction with hand cutting where divers dislodge the plants. The plant/sediment slurry is hydraulically pumped to the barge through hoses carried by the diver. Its effectiveness is dependent on sediment composition, density of aquatic plants, and underwater visibility. Suction dredging may be best suited for localized infestations of low plant density where fragmentation must be controlled. A permit will be required for this activity.

Advantages: Diver suction dredging is species –selective. Disruption of sediments can be minimized. These methods can remove the plant stems and roots, resulting in thorough plant disruption and potential longer term control. Fragmentation of plants is minimized. This activity can be completed near and around obstacles such as piers or marinas where a harvester could not operate.

Disadvantages: Diver suction dredging is labor intensive and costly. Upland disposal of dredged slurry can require additional equipment and costs. Increased turbidity in the area of treatment can be a problem. Release of nutrients and other pollutants can also be a problem.

Costs: Suction dredging costs can be variable depending on equipment and transport requirements for slurry. Costs range from \$5,000 per acre to \$10,000 per acre.

Dredging

Sediment removal through dredging can work as a plant control technique by limiting light through increased water depth or removing soft sediments that are a preferred habitat to nuisance rooted plants. Soft sediment removal is accomplished with drag lines, bucket dredges, long reach backhoes, or other specialized dredging equipment. Dredging has had mixed results in controlling aquatic plant, however it can be highly effective in appropriate situations. Dredging is most often applied in a major restructuring of a severely degraded system. Generally, dredging is an activity associated with other restoration efforts. Comprehensive pre-planning will be necessary for these techniques and a dredging permit would be required.

Advantages: Dredging can remove nutrient reserves which result in nuisance rooted aquatic plant growth. Dredging, when completed, can also actually improve substrate and habitat for more desirable species of aquatic plants, fish, and invertebrates. It allows the complete renovation of an aquatic ecosystem. This method has the potential for significant reductions in aquatic plant growth. These methods can be completed in “off season” months avoiding interference with summer recreation activities.

Disadvantages: Dredging can temporarily destroy important fish and invertebrate habitat. Suspension of sediments usually increases turbidity significantly and can possibly release nutrients causing algae blooms. Dredging is extremely expensive and requires significant planning. Dredged materials may contain toxic materials (metals, PCBs). Dredged material transportation and disposal of toxic materials are additional management considerations and are potentially expensive. It could be difficult and costly to secure regulatory permits and approvals.

Costs: Dredging costs depend upon the scale of the project and many other factors. It is generally an extremely expensive option.

Drawdown: Water level drawdown exposes the plants and root systems to prolonged freezing and drying to kill the plants. It can be completed any time of the year, however is generally more effective in winter, exposing the lake bed to freezing temperatures. If there is a water level control structure capable of drawdown, it can be an in-expensive way to control some aquatic plants. Aquatic plants vary in their susceptibility to drawdown, therefore, accurate identification of problem species is important. Drawdown is often used for other purposes of improving waterfowl habitat or fishery management, but sometimes has the added benefit of nuisance rooted aquatic plant control. This method can be used in conjunction with a dredging project to excavate nutrient-rich sediments. This method is best suited for use on reservoirs or shallow man-made lakes. A drawdown would require regulatory permits and approvals.

Advantages: A drawdown can result in compaction of certain types of sediments and can be used to facilitate other lake management activities such as dam repair, bottom barrier, or dredging projects. Drawdown can significantly impact populations of aquatic plants that propagate vegetatively. It is inexpensive.

Disadvantages: This method is limited to situations with a water level control structure. Pumps can be used to de-water further if groundwater seepage is not significant. This technique may also result in the removal of beneficial plant species. Drawdowns can decrease bottom dwelling invertebrates and overwintering reptiles and amphibians. Drawdowns can affect adjacent wetlands, alter downstream flows, and potentially impair well production. Drawdowns and any water level manipulation are often highly controversial since shoreline landowners access and public recreation are limited during the drawdown. Fish populations are vulnerable during a drawdown due to over-harvesting by fisherman in decreased water volumes.

Costs: If a suitable outlet structure is available then costs should be minimal. If dewatering pumps would be required or additional management projects such as dredging are completed, additional costs would be incurred. Other costs would include recreational losses and perhaps loss in tourism revenue.

Chemical Controls

Using chemical herbicides to kill nuisance aquatic plants is the oldest APM method. However, past pesticides uses being linked to environmental or human health problems have led to public wariness of chemicals in the environment. Current pesticide registration procedures are more stringent than in the past. While no chemical pesticide can be considered 100 percent safe, federal pesticide regulations are based on the premise that if a chemical is used according to its label instructions it will not cause adverse environmental or human health effects.

Chemical herbicides for aquatic plants can be divided into two categories, systemic and contact herbicides. Systemic herbicides are absorbed by the plant, translocated throughout the plant, and are capable of killing the entire plant, including the roots and shoots. Contact herbicides kill the plant surface in which it comes in contact, leaving roots capable of re-growth. Aquatic herbicides exist under various trade names, causing some confusion. Aquatic herbicides include the following:

- ▲ Endothall Based Herbicide
- ▲ Diquat Based Herbicide
- ▲ Fluridone Based Herbicide
- ▲ 2-4 D Based Herbicide
- ▲ Glyphosate Based Herbicide
- ▲ Triclopyr Based Herbicide
- ▲ Phosphorus Precipitation

Each of these methods are described below. The costs, benefits, and drawbacks of each chemical APM alternative are provided.

Endothall Based Herbicide: Endothall is a contact herbicide, attacking a wide range of plants at the point of contact. The chemical is not readily transferred to other plant tissue, therefore regrowth can be expected and repeated treatments may be needed. It is sold in liquid and granular forms under the trade names of Aquathol[®] or Hydrothol[®]. Hydrothol is also an algaecide. Most endothall products break down easily and do not remain in the aquatic environment. Endothall products can result in plant reductions for a few weeks to several months. Multi-season effectiveness is not typical. A permit is required for use of this herbicide.

Advantages: Endothall products work quickly and exhibit moderate to highly effective control of floating and submersed species. This herbicide has limited toxicity to fish at recommended doses.

Disadvantages: The entire plant is not killed when using endothall. Endothall is non-selective in the treatment area. High concentrations can kill fish easily. Water use restrictions (time delays) are necessary for recreation, irrigation, and fish consumption after application.

Costs: Costs vary with treatment area and dosage. Average costs for chemical application range between \$400 and \$700 per acre.

Diquat Based Herbicide: Diquat is a fast-acting contact herbicide effective on a broad spectrum of aquatic plants. It is sold under the trade name Reward[®]. Diluted forms of this product are also sold as private label products. Since Diquat binds to sediments readily, its effectiveness is reduced by turbid water. Multi-season effectiveness is not typical. A permit is required for use of this herbicide.

Advantages: Diquat works quickly and exhibit moderate to highly effective control of floating and submersed species. This herbicide has limited toxicity to fish at recommended doses.

Disadvantages: The entire plant is not killed when using diquat. Diquat is non-selective in the treatment area. Diquat can be inactivated by suspended sediments. Diquat is sometimes toxic to zooplankton at the recommended dose. Limited water used restrictions (water supply, agriculture, and contact recreation) are required after application.

Costs: Costs vary with treatment area and dosage. A general cost estimate for treatment is between \$200 and \$500 per acre.

Fluoridone Based Herbicide: Fluoridone is a slow-acting systemic herbicide, which is effectively absorbed and translocated by both plant roots and stems. Sonar[®] and Avast![®] is the trade name and it is sold in liquid or granular form. Fluoridone requires a longer contact time and demonstrates delayed toxicity to target plants. Eurasian watermilfoil is more sensitive to fluoridone than other aquatic plants. This allows a semi-selective approach when low enough doses are used. Since the roots are also killed, multi-season effectiveness can be achieved. It is best applied during the early growth phase of the plants. A permit and extensive planning is required for use of this herbicide.

Advantages: Fluoridone is capable of killing roots, therefore producing a longer lasting effect than other herbicides. A variety of emergent and submersed aquatics are susceptible to this herbicide. Fluoridone can be used selectively, based on concentration. A gradual killing of target plants limits severe oxygen depletion from dead plant material. It has demonstrated low toxicity to aquatic fauna such as fish and invertebrates. 3 to 5 year control has been demonstrated. Extensive testing has shown that, when used according to label instructions, it does not pose negative health affects.

Disadvantages: Fluoridone is a very slow-acting herbicide sometimes taking up to several months for visible effects. It requires a long contact time. Fluoridone is extremely soluble and mixable, therefore, not effective in flowing water situations or for treating a select area in a large open lake. Impacts on non-target plants are possible at higher doses. Time delays are necessary on use of the water (water supply, irrigation, and contact recreation) after application.

Costs: Costs vary with treatment area and dosage. Treatment costs range from \$500 to \$2,000 per acre.

2,4-D Based Herbicide: 2,4-D based herbicides are sold in liquid or granular forms under various trade names. Common granular forms are sold under the trade names Navigate[®] and Aqua Kleen[®]. Common liquid forms include DMA 4[®] and Weedar 64[®]. 2,4-D is a systemic herbicide that affects broad leaf plants. It has been demonstrated effective against Eurasian watermilfoil, but it may not work on many aquatic plants. Since the roots are also killed, multi-season effectiveness may be achieved. It is best applied during the early growth phase of the plants. Visible results are evident within 10 to 14 days. A permit is required for use of this herbicide.

Advantages: 2,4-D is capable of killing roots, therefore producing a longer lasting effect than some other herbicides. It is fairly fast and somewhat selective, based on application timing and concentration. 2,4-D containing products are moderately to highly effective on a few emergent, floating, or submersed plants.

Disadvantages: 2,4-D can have variable toxicity effects to aquatic fauna, depending on formulation and water chemistry. 2,4-D lasts only a short time in water, but can be detected in sediments for months after application. Time delays are necessary on use of the water (agriculture and contact recreation) after application. The label does not permit use of this product in water used for drinking, irrigation, or livestock watering.

Costs: Costs vary with treatment area and dosage. Treatment costs range from \$300 to \$800 per acre.

Glyphosate Based Herbicide: Glyphosate has been categorized as both a contact and a systemic herbicide. It is applied as a liquid spray and is sold under the trade name Rodeo[®] or Pondmaster[®]. It is a non-selective, broad based herbicide effective against emergent or floating leaved plants, but not submergents. It's effectiveness can be reduced by rain. A permit is required for use of this herbicide.

Advantages: Glyphoshate is moderately to highly effective against emergent and floating-leaf plants resulting in rapid plant destruction. Since it is applied by spraying plants above the surface, the applicator can apply it selectively to target plants. Glyphosate dissipates quickly from natural waters, has a low toxicity to aquatic fauna, and carries no restrictions or time delays for swimming, fishing, or irrigation.

Disadvantages: Glyphoshate is non-selective in the treatment area. Wind can dissipate the product during the application reducing it's effectiveness and cause damage to non-target organisms. Therefore, spray application should only be completed when wind drift is not a problem. This compound is highly corrosive, therefore storage precautions are necessary.

Costs: Costs average \$500 to \$1,000 per acre depending on the scale of treatment.

Triclopyr Based Herbicide: Triclopyr is a systemic herbicide. It is registered for experimental aquatic use in selected areas only. It is applied as a liquid spray or injected into the subsurface as a liquid. Triclopyr is sold under the trade name Renovate[®] or Restorate[®]. Triclopyr has shown to be an effective control to many floating and submersed plants. It has been demonstrated to be highly effective against Eurasian watermilfoil, having little effect on valued native plants such as pondweeds. Triclopyr is most effective when applied during the active growth period of younger plants.

Advantages: This herbicide is fast acting. Triclopyr can be used selectively since it appears more effective against dicot plant species, including several difficult nuisance plants. Testing has demonstrated low toxicity to aquatic fauna.

Disadvantages: At higher doses, there are possible impacts to non-target species. Some forms of this herbicide are experimental for aquatic use and restrictions on use of the treated water are not yet certain.

Biological Controls

There has been recent interest in using biological technologies to control aquatic plants. This concept stems from a desire to use a “natural” control and reduce expenses related to equipment and/or chemicals. While use of biological controls is in its infancy, potentially useful technologies have been identified and show promise for integration with physical and chemical APM strategies. Several biological controls that are in use or are under experimentation include the following:

- ▲ Herbivorous Fish
- ▲ Herbivorous Insects
- ▲ Plant Pathogens
- ▲ Native Plants

Each of these methods are described below. The costs, benefits, and drawbacks of each biologic APM method are provided.

Herbivorous Fish: A herbivorous fish such as the non-native grass carp can consume large quantities of aquatic plants. These fish have high growth rates and a wide range of plant food preferences. Stocking rates and effectiveness will depend on many factors including climate, water temperature, type and extent of aquatic plants, and other site-specific issues. Sterile (triploid) fish have been developed resulting in no reproduction of the grass carp and population control. This technology has demonstrated mixed results and is most appropriately used for lake-wide, low intensity control of submersed plants. Some states do not allow stocking of herbivorous fish. In Wisconsin, stocking of grass carp is prohibited.

Advantages: This technology can provide multiple years of aquatic plant control from a single stocking. Compared to other long-term aquatic plant control techniques such as bottom tillage or bottom barriers, costs may be relatively low.

Disadvantages: Sterile grass carp exhibit distinct food preferences, limiting their applicability. Grass carp may feed selectively on the preferred plants, while less preferred plants, including milfoil, may increase. The effects of using grass carp may not be immediate. Overstocking may result in an impact on non-target plants or eradication of beneficial plants, altering lake habitat. Using grass carp may result in algae blooms and increased turbidity. If precautions are not taken (i.e. inlet and outlet control structures to prevent fish migration) the fish may migrate and have adverse effects on non-target vegetation.

Costs: Costs can range from \$50/acre to over \$2,000/acre, at stocking rates of 5 fish/acre to 200 fish/acre.

Herbivorous Insects: Non-native and native insect species have been used to control rooted plants. Using herbivorous insects is intended to selectively control target species. These aquatic larvae of moths, beetles, and thrips use specific host aquatic plants. Several non-native species have been imported under USDA approval and used in integrated pest management programs, a combination of biological, chemical, and mechanical controls.

These non-native insects are being used in southern states to control nuisance plant species and appear climate-limited, their northern range being Georgia and North Carolina. While successes have been demonstrated, non-native species have not established themselves for solving biological problems, sometimes creating as many problems as they solve. Therefore, government agencies prefer alternative controls.

Native insects such as the larvae of midgeflies, caddisflies, beetles, and moths may be successful APM controls in northern states. Recently however, the native aquatic weevil *Euhrychiopsis lecontei* has received the most attention. This weevil has been associated with native northern water milfoil. The weevil can switch plant hosts and feed on Eurasian watermilfoil, destroying its growth points. While the milfoil weevil is gaining popularity, it is still experimental.

Advantages: Herbivorous insects are expected to have no negative effects on non-target species. The insects have shown promise for long term control when used as part of integrated aquatic plant management programs. The milfoil weevils do not use non-milfoil plants as hosts.

Disadvantages: Natural predator prey cycles indicate that incomplete control is likely. An oscillating cycle of control and re-growth is more likely. Fish predation may complicate controls. Large numbers of milfoil weevils may be required for a dense stand and can be expensive. The weevil leaves the water during the winter, may not return to the water in the spring, and are subject to bird predation in their terrestrial habitat. Application is manual and extremely time consuming. Introducing any species, especially non-native ones, into an aquatic ecosystem may have undesirable effects. Therefore, it is extremely important to understand the life cycles of the insects and the host plants.

Costs: Reported costs of herbivorous insects rang from \$300/acre to \$3,000/acre.

Specifically, the native milfoil weevils cost approximately \$1.00 per weevil. It is generally considered appropriate to use 5 to 7 weevils per stem. Dense stands of milfoil may contain 1 to 2 million stems per acre. Therefore, costs of this new technology are currently prohibitive.

Plant Pathogens: Using a plant pathogen to control nuisance aquatic plants has been studied for many years, however, plant pathogens still remain largely experimental. Fungi are the most common pathogens, while bacteria and viruses have also been used. There is potential for highly specific plant applications.

Advantages: Plant pathogens may be highly species specific. They may provide substantial control of a nuisance species.

Disadvantages: Pathogens are experimental. The effectiveness and longevity of control is not well understood. Possible side effects are also unknown.

Costs: These techniques are experimental therefore a supply of specific products and costs are not established.

Native Plants: This method involves removing the nuisance plant species through chemical or physical means and re-introducing seeds, cuttings, or whole plants of desirable species. Success has been variable. When using seeds, they need to be planted early enough to encourage the full growth and subsequent seed production of those plants. Transplanting mature plants may be a better way to establish seed producing populations of desirable aquatics. Recognizing that a healthy, native, desirable plant community may be resistant to infestations of nuisance species, planting native plants should be encouraged as an APM alternative. Non-native plants can not be translocated.

Advantages: This alternative can restore native plant communities. It can be used to supplement other methods and potentially prevent future needs for costly repeat APM treatments.

Disadvantages: While this appears to be a desirable practice, it is experimental at this time and there are not many well documented successes. Nuisance species may eventually again invade the areas of native plantings. Careful planning is required to ensure that the introduced species do not themselves become nuisances. Hand planting aquatic plants is labor intensive.

Costs: Costs can be highly variable depending on the selected native species, numbers of plants ordered, and the nearest dealer location.

Aquatic Plant Prevention

The phrase “an ounce of prevention is worth a pound of cure” certainly holds true for APM. Prevention is the best way to avoid nuisance aquatic plant growth. Prevention of the spread of invasive aquatic plants must also be achieved. Inspecting boats, trailers, and live wells for live aquatic plant material is the best way to prevent nuisance aquatic plants from entering a new aquatic ecosystem. Protecting the desirable native plant communities is also important in maintaining a healthy aquatic ecosystem and preventing the spread of nuisance aquatics once they are present.

Prolific growth of nuisance aquatic plants can be prevented by limiting nutrient (i.e. phosphorus) inputs to the water body. Aeration or phosphorus precipitation can achieve controls of in-lake cycling of phosphorus, however, if there are additional outside sources of nutrients, these methods will be largely ineffective in controlling algae blooms or intense aquatic macrophyte infestations. Watershed management activities to control nutrient laden storm water runoff are critical to controlling excessive nutrient loading to the water bodies. Nutrient loading can be prevented/minimized by the following:

- ▲ Shoreline buffers
- ▲ Using non-phosphorus fertilizers on lawns
- ▲ Settling basins for storm water effluents