



Report

Aquatic Plant Management Plan

Crab Lake

Presque Isle Town Lakes Committee

September 30, 2011

Project Number 004010-09001-0

WDNR Grant LPL-1135-07

1203 Storbeck Drive
Waupun, WI 53963

Tel 920-324-8600
Fax 920-324-3023

www.bonestroo.com



AQUATIC PLANT MANAGEMENT PLAN

CRAB LAKE

September 30, 2011

Prepared For:

Presque Isle Town Lakes Committee
7595 Annabelle Lake Road
Presque Isle, Wisconsin 54557

Prepared By:

Bonestroo
1203 Storbeck Drive
Waupun, Wisconsin 53963

Project Number: 004010-09001-0

A handwritten signature in blue ink, appearing to read "James T. Scharl", written over a horizontal line.

James T. Scharl
Graduate Scientist

Table of Contents

1.0 Executive Summary	1
2.0 Introduction.....	3
3.0 Baseline Information	4
3.1 Lake History and Morphology.....	4
3.2 Watershed Overview.....	4
3.3 Water Quality.....	4
3.3.1 Water Clarity	5
3.3.2 Total Phosphorus and Chlorophyll a	5
3.3.3 Dissolved Oxygen and Temperature	7
3.3.4 Trophic State Index.....	7
3.4 Summary of Lake Fishery	8
3.5 Lake Management History	9
3.6 Goal and Objectives.....	9
4.0 Project Methods	10
4.1 Existing Data Review	10
4.2 Aquatic Plant Survey and Analysis	10
4.3 Shoreline Characterization	11
4.4 Public Involvement, Questionnaire, and Plan Review	12
5.0 Discussion of Project Results	13
5.1 Aquatic Plant Ecology.....	13
5.2 Aquatic Invasive Species	13
5.3 2007 Aquatic Plant Survey.....	13
5.3.1 Floating-Leaf Plants.....	15
5.3.2 Submergent Plants.....	15
5.3.3 Emergent Plants	16
5.4 North Crab Lake Plant Survey	16
5.4.1 Floating-Leaf Plants.....	16
5.4.2 Submergent Plants.....	16
5.4.3 Emergent Plants	17
5.5 Floristic Quality Index	17
5.6 Shoreline Characterization	17
5.7 Public Questionnaire	17
5.8 Water Quality Sampling.....	18
6.0 Management Alternatives and Recommendations.....	20
6.1 Aquatic Plant Maintenance Alternatives.....	20
6.1.1 Aquatic Invasive Species.....	20
6.1.2 Clean Boats/Clean Waters Campaign	21
6.1.3 Aquatic Plant Protection and Shoreline Management	21

6.1.4 Public Education and Involvement	22
6.2 Aquatic Plant Manipulation Alternatives	22
6.2.1 Manual Removal	22
6.2.2 Aquatic Invasive Plant Species Chemical Herbicide Treatment	23
7.0 Conclusion and Recommended Action Plan	25
7.1 Recommended Active Goals.....	25
7.2 Closing	26
8.0 References	27

FIGURES

Lake Location and Watershed	1
Bathymetric Map	2
Point Intercept Sampling Locations	3
August 2007 Aquatic Plant Distribution.....	4a – 4g
Shoreline Characterization	5
Point Sample Locations and Shoreline Characterization – North Crab Lake	6

TABLES

Taxa Detected During 2007 Aquatic Plant Survey	1
2007 Aquatic Plant Community Statistics	2
2007 Aquatic Plant Taxa-Specific Statistics	3
Floristic Quality Index	4
Taxa Detected During 2007 Aquatic Plant Survey – North Crab Lake.....	5
Floristic Quality Index – North Crab Lake.....	6

APPENDIX

Aquatic Plant Management Tables, Figures and Point Intercept Sample Coordinates	A
Summary of Public Survey	B
Importance of Aquatic Plants to Lake Ecosystem.....	C1
Aquatic Invasive Species.....	C2
Descriptions of Aquatic Plants	D
Summary of Aquatic Plant Management Alternatives	E
NR 107 and NR109 Wisconsin Administrative Code	F
Resource for Additional Information.....	G
Aquatic Plant Management Strategy, Northern Region WDNR	H
Temperature and Dissolved Oxygen Profiles	I
Additional Information / Lake Resident Comments	J

1.0 Executive Summary

The Presque Isle Town Lakes Committee (PITLC) was formed in 2005 to address resource management concerns in lakes within the Town of Presque Isle. The Committee has been active in a number of lake management activities on Crab Lake including: aquatic plant management, water quality sampling, invasive species sampling, and community education activities. The Committee contracted Northern Environmental (now Bonestroo, Inc.) to help develop an aquatic plant management (APM) plan for Crab Lake. The Crab Lake APM Plan includes a review of available lake information, an aquatic plant survey, and an evaluation of feasible physical, mechanical, biological, and chemical management alternatives if deemed appropriate. The APM plan also recommends specific prevention activities for aquatic invasive species (AIS) in the lake system; which are discussed below.

Northern Environmental completed an aquatic plant survey on Crab Lake in 2007, which identified twenty seven aquatic plant species. The most abundant aquatic plants identified during the survey were wild celery (*Vallisneria americana*), dwarf watermilfoil (*Myriophyllum tenellum*), and three equally abundant species creeping spikerush (*Eleocharis palustris*), large-leaf pondweed (*Potamogeton amplifolius*), and fern pondweed (*Potamogeton robbinsii*). A brief aquatic plant survey was also conducted on North Crab Lake identifying twenty seven aquatic plant species. The Floristic Quality Index (FQI) is an index that uses the aquatic plant community as an indicator of lake health. Crab Lake exhibited an FQI of 37.07, higher than the state northern ecoregion average (24.3). North Crab Lake exhibited and FQI of 34.1, also higher than the northern ecoregion average.

RECOMMENDED AQUATIC PLANT MANAGEMENT PLAN

No aquatic invasive plants were found during the aquatic plant survey in 2007. If an invasive plant were found the Wisconsin Department of Natural Resources (WDNR) recommends that Crab Lakes residents work with WDNR and aquatic plant professionals to determine extent of AIS after discovery and to then determine appropriate management (per WDNR Water Resource Management Specialist's, Kyle McLaughlin, comments dated December 22, 2008). The fact that the native plant community has an above average FQI score illustrates that there is a unique and diverse plant community in Crab Lake. Such a plant community is worthy of protection from human disturbance and from the impact aquatic invasive species would have if introduced to this system. Because of that, the following Recommended Action Plan focuses on conservation and plant protection.

The following Active Goals form the structure of the Crab Lake Aquatic Plant Management Plan:

Active Goal: To continue and expand the WDNR Clean Boats, Clean Waters program on Crab Lake.

Active Goal: To provide visitors with educational information concerning the potential impact their activities could have on introduction of aquatic invasive species, wildlife, habitats and Crab Lake water quality.

- Active Goal:** To implement and maintain an aquatic invasive species monitoring program that will survey for invasive species, and if found, monitor their locations and extent of population spread.
- Active Goal:** To continue the Crab Lake comprehensive water quality monitoring program through the WDNR Citizen Lake Monitoring Network
- Active Goal:** To work in concert with the WDNR staff and representatives of fishing related businesses to evaluate Crab Lake fish management practices and develop goals in order to maintain and enhance a quality family sport fishery.
- Active Goal:** To develop and implement a fishing tournament AIS management and monitoring program in order to be prepared if fishing tournaments come to Crab Lake.
- Active Goal:** To enforce the Town of Presque Isle's 200 foot no-wake areas (from shoreline and islands) ordinances in order to minimize recreational impacts on the plant community, shoreline habitats, and to promote safe boating.
- Active Goal:** To support the identification and preservation of critical species and critical habitat lands, and wetlands within the watershed. (These are areas with rare vegetation, important habitat for wildlife, or important spawning and nursery areas for fish. Preservation of these lands has a direct impact on the water quality of the lake).
- Active Goal:** To provide education and information to shoreline property owners regarding how native aquatic plant protection and shoreline management can slow the spread of aquatic invasive plants (if they become introduced), improve the lake fishery, improve wildlife habitat and affect the quality of the water in the lake (including development of a *shoreline restoration packet* that could be given to landowners who's property has development categorized as Moderate or Major).
- Active Goal:** To encourage the incorporation of water quality protection measures in the design, construction and maintenance of all lake access sites on Crab Lake (e.g. storm water control, site drainage control, appropriate plant matter disposal, and watercraft wash down facilities if found to be needed). (*However, see Appendix J.*)
- Active Goal:** To meet on a regular basis with local government agencies and representatives of lakes located within the Town of Presque Isle, to identify essential and new lake management issues and determine collaborative solutions.

2.0 Introduction

Crab Lake is a 949 acre drainage lake located in northern Vilas County. North Crab Lake is a 56 acre drainage lake connected by a navigable channel to Crab Lake. The two lakes have a 12,290 acre (19.2 square miles) watershed. Crab Lake exhibits good water clarity and according to the Wisconsin Trophic State Index, is a mesotrophic lake.

Lake residents have become concerned about the possibility of the introduction of Eurasian watermilfoil (*Myriophyllum spicatum* - EWM), curly-leaf pondweed (*Potamogeton crispus* - CLP) and other AIS into the aquatic plant community of Crab Lake. Although no AIS were recorded during the aquatic plant survey in 2007, this APM Plan includes strategies for detection, monitoring, and management/removal of EWM and CLP from Crab Lake if ever established.

This document is the APM Plan for Crab Lake and discusses the following:

- Lake morphology and lake watershed characteristics
- Historical aquatic plant management activities
- Stakeholder's goals and objectives
- Aquatic plant ecology
- 2007 baseline aquatic plant survey
- Feasible aquatic plant management alternatives
- Selected suite of aquatic plant management options

3.0 Baseline Information

3.1 LAKE HISTORY AND MORPHOLOGY

Crab Lake is located in the Town of Presque Isle in the northern Vilas County, Wisconsin. The lake is part of the headwaters area of the Presque Isle-Black River Watershed, which drains north to Lake Superior. Figure 1 depicts the lake location. The following summarizes the lake's physical attributes:

Lake Name	Crab	North Crab
Lake Type	Drainage	Drainage
Surface Area (acres)	949	56
Maximum depth (feet)	64	35
Mean depth (feet)	22	NA
Shoreline Length (miles)	15.76	NA
Public Landing	Yes	No

Source: Wisconsin Lakes, WDNR 2005 and WDNR Lake Survey map, 1941

Figure 2 illustrates the lake bathymetries. Crab Lake provides year-round recreation activities ranging from, fishing, swimming, waterskiing, pleasure boating, snowmobiling, and more.

3.2 WATERSHED OVERVIEW

The Crab Lake watershed encompasses approximately 12,290 acres in northern Vilas County and is primarily forested and undeveloped. Crab Lake is connected by small waterways to the Aimer Lakes, Knife Lake, Annabelle Lake, Belle Lake, Crab Lake, North Crab Lake, Armour Lake, and Horsehead Lake before draining to the Presque Isle River. Crab Lake and North Crab Lake are listed as a Priority Navigable Water (PNW) on the Wisconsin list of Designated Waters. Priority Navigable Waters include waters with self-sustaining walleye populations in ceded territories and waters with self-sustaining musky populations. Crab Lake contains both a self-sustaining walleye population and a self-sustaining muskellunge population within the Ceded Territories.

The Crab Lake area consists mainly of Gogebic and Fence soil types. These soils are well drained, sandy soils formed in glacial till and glacial lake plains with stratified silty and sandy loam deposits. They generally have slopes of 0 to 15 percent. A fragipan layer sometimes occurs within Gogebic soils. Fragipans are loamy, brittle subsurface horizons consisting of clay, silt and sand that appear cemented and can restrict plant root growth (USDA, 1988).

3.3 WATER QUALITY

WDNR Lake Water Quality Database indicates that the following water quality information is available


- Water clarity (Secchi depth) 1992 - 2007 (Citizen Lake Monitoring)
- Total phosphorus – 1993 - 2007 (Citizen Lake Monitoring)
- Chlorophyll a – 1993 - 2007 (Citizen Lake Monitoring)
- Temperature & dissolved oxygen profiles – 1993-2007 (Citizen Lake Monitoring)

The above referenced data was used in creating the Crab APM Plan. Higher Secchi depth readings indicate clearer water and deeper light penetration. Total Phosphorus is a measure of nutrients available for plant growth. Chlorophyll a is green pigment present in all plant life and necessary for photosynthesis. The amount present in lake water depends on the amount of algae suspended in the water column of a lake. Chlorophyll a is used as a common indicator of water quality (Shaw et al, 2004). Higher chlorophyll a values indicate lower water qualities

3.3.1 WATER CLARITY


The historical water clarity average based on Secchi Disk readings is 13.85 feet (2.28 meters) and ranges from eight to 18 feet. The Wisconsin average Secchi Disk reading in 2005 was 10 feet (Larry Bresina, The Secchi Disk and Our Eyes - Working Together To Measure Clarity of Our Lakes; internet document). The following graph illustrates the historical water clarity measurements on Crab Lake, data collected in 2007 is included.

Crab Lake Secchi Readings




3.3.2 TOTAL PHOSPHORUS AND CHLOROPHYLL A

Historically Crab Lake has an average phosphorus reading of 0.0114 milligrams per liter (parts per million – ppm). Total phosphorus has varied from 1993 through 2007 from 0.006 ppm to 0.029 ppm. The following graph illustrates the historical phosphorus measurements on Crab Lake.



Chlorophyll measurements started in 1993 and continued through 2007, showing an average of 2.54 micrograms per liter (parts per billion – ppb). Data ranged from 1.00 ppb to 4.99 ppb. The following graph illustrates the historical chlorophyll measurements on Crab Lake.



3.3.3 DISSOLVED OXYGEN AND TEMPERATURE

WDNR data indicates that profile data was collected over the deepest portion of the lake for dissolved oxygen (DO) and temperature from 1993-2007 (Appendix I). Dissolved oxygen data was not available for 1993-1994.

3.3.4 TROPHIC STATE INDEX

Trophic State Index (TSI) values are assigned to a lake based on total phosphorus, chlorophyll *a*, and water clarity values. The TSI is a measure of a lake's biological productivity. The TSI used for Wisconsin lakes is described below.

Category	TSI	Lake Characteristics	Total P (ug/l)	Chlorophyll a (ug/l)	Water Clarity (feet)
Oligotrophic	1-40	Clear water; oxygen rich at all depths, except if close to mesotrophic border; then may have low or no oxygen; cold-water fish likely in deeper lakes.	< 12	<2.6	>13
Mesotrophic	41-50	Moderately clear; increasing probability of low to no oxygen in bottom waters.	12 to 24	2.6 to 7.3	13 to 6.5
Eutrophic	51-70	Decreased water clarity; probably no oxygen in bottom waters during summer; warm-water fisheries only; blue-green algae likely in summer in upper range; plants also excessive.	> 24	>7	<6.5
Crab Lake	39.41	Oligotrophic	11.4	2.54	13.85

Adopted from Carlson 1977, Lillie and Mason, 1983, and Shaw 1994 et. al.

The historical water clarity, total phosphorus, and chlorophyll *a* data indicate that Crab Lake is an oligotrophic lake. Data collected during by Citizen Lake Monitors is included in the above averages.

3.4 SUMMARY OF LAKE FISHERY

The following table identifies the fish species the WDNR lists as being present in Crab Lake.

Fish Species	Present	Common	Abundant
Muskellunge		X	
Northern Pike	X		
Walleye		X	
Largemouth Bass	X		
Smallmouth Bass	X		
Panfish		X	

Source: WDNR Wisconsin Lakes Publication # PUB-FH-800, 2005

The WDNR website presents records of fish stocked in Crab Lake (WDNR Fish stocking website, 2007). All walleye were stocked as fingerling while muskellunge have been stocked as fingerlings in 1980-1996, large fingerlings in 1998-2006, and yearlings in 1990.

Year	Musky	Walleye
1974		10200
1979		47040
1980	1041	
1984	759	
1985	1065	
1988	1000	
1990	1000*	
1991	400	
1992	500	
1996	450	
1998	900	
2000	900	
2002	473	
2004	474	
2006	474	

* Both fingerlings (800) and yearlings (200) stocked

All fisheries in Crab Lake are currently sustained through natural reproduction. Crab Lake is located in the "Ceded Territories" of Wisconsin. The Ceded Territories was ceded to the United States by the Lake Superior Chippewa Tribes in 1837 and 1842. The WDNR describes Native American fishing in the Ceded Territories this way: "The six Chippewa tribes of Wisconsin are legally able to harvest walleyes using a variety of high efficiency methods, but spring spearing is the most frequently used method. In spring each tribe declares how many walleyes and muskellunge they intend to harvest from each lake. Harvest begins shortly after ice-out, with nightly fishing permits issued to individual tribal spearers. Each permit allows a specific number of fish to be harvested, including one walleye between 20 and 24 inches and one additional walleye of any size. All fish that are taken are documented each night with a tribal clerk or

warden present at each boat landing used in a given lake. Once the declared harvest is reached in a given lake, no more permits are issued for that lake and spearfishing ceases (<http://dnr.wi.gov/fish/ceded/tribalharvest.html>)."

All fishing regulations and bag limits for Crab are concurrent with standard WDNR regulations in the Ceded Territories except for walleye and muskellunge. For non-Chippewa anglers there is no minimum size limit for walleye with a three (3) fish daily bag limit, but only one may be over 15 inches. In the Ceded Territories the WDNR works to establish "safe harvest limits" set so there is less than a 1-in-40 chance that more than 35% of the adult walleye population will be harvested in any given lake by either tribal or recreational fishermen, or both combined. For muskellunge, the daily bag limit is one with a minimum size of 40 inches.

3.5 LAKE MANAGEMENT HISTORY

The only past management activities on Crab Lake are fish stocking by the WDNR. The historical WDNR stocking data is presented in the previous section. Overall, the Town of Presque Isle has adopted several ordinances that help protect lakes and shorelines. Town of Presque Isle ordinances 500, 902, 908, and 909 all protect lakes in one way or another. Ordinance 501.03(2) mandates no wake on lakes less than 50 acres. 501.03(3),(a) requires no wake within 200 feet of any shoreline. 501.03(4) limits waterskiing and other like activities to the hours between 10 AM and 5 PM. The above mentioned ordinances in the 900 category (902.1.1), zoning, limit lot sizes within 600 feet of Crab Lake to 400 feet of frontage and 5 acres on all lots created after May 8, 1959. Other watershed related ordinances are included in the zoning ordinances.

3.6 GOALS AND OBJECTIVES

PITLC identified the following goals for aquatic plant management on Crab Lake.

- Maintain and improve recreational opportunities
- Protect and improve fish and wildlife habitat
- Preserve native aquatic plants
- Prevent the introductions of AIS
- Identify and Protect sensitive areas
- Identify sources of financial assistance for aquatic plant management activities
- Coordinate sound aquatic plant management practices where needed within Crab Lake
- Educate the Crab Lake community on proper AIS identification and prevention efforts
- Gather citizen input
- Increase citizen participation in lake management

4.0 Project Methods

To accomplish the project goals, the PITLC needs to make informed decisions regarding APM on the lake. To make informed decisions, PITLC proposed to:

- Collect, analyze, and interpret basic aquatic plant community data
- Recommend practical, scientifically-sound aquatic plant management strategies

Offsite and onsite research methods were used during this study. Offsite methods included a thorough review of available background information on the lake, its watershed, and water quality. An aquatic plant community survey was completed onsite to provide the data needed to evaluate aquatic plant management alternatives.

4.1 EXISTING DATA REVIEW

Bonestroo researched a variety of information resources to develop a thorough understanding of the ecology of the Lake. Information sources included:

- Local and regional geologic, limnologic, hydrologic, and hydrogeologic research
- Discussions with lake members
- Available topographic maps and aerial photographs
- Data from WDNR files

These sources were essential to understanding the historic, present, and potential future conditions of the lake, as well as to ensure that previously completed studies were not unintentionally duplicated. Specific references are listed in Section 8.0 of this report.

4.2 AQUATIC PLANT SURVEY AND ANALYSIS

The aquatic plant community of the lake was surveyed on August 2 - 4, 2007 by Northern Environmental. The survey was completed according to the point intercept sampling method described by Madsen (1999) and as outlined in the WDNR draft guidance entitled "Aquatic Plant Management in Wisconsin" (WDNR, 2005).

WDNR research staff determined the sampling point resolution in accordance with the WDNR guidance and provided a base map with the specified sample point locations. The sample resolution was a 55 meter grid with 1204 pre-determined intercept points (Figure 3). Latitude and longitude coordinates and sample identifications were assigned to each intercept point on the grid (Appendix A). Geographic coordinates were uploaded into a global positioning system (GPS) receiver. The GPS unit was then used to navigate to intercept points. At each intercept point, plants were collected by tossing a specialized rake on a rope and dragging the rake along the bottom sediments. All collected plants were identified to the lowest practicable taxonomic level (e.g., typically genus or species) and recorded on field data sheets. Visual observations of aquatic plants were also recorded. Water depth and, when detectable, sediment types at each intercept point were also recorded on field data sheets.

The point intercept method was used to evaluate the existing emergent, submergent, floating-leaf, and free-floating aquatic plants. If a species was not collected at a specific point, the space

on the datasheet was left blank. For the survey, the data for each sample point was entered into the WDNR “Worksheets” (i.e., a data-processing spreadsheet) to calculate the following statistics:

- **Taxonomic richness** (the total number of taxa detected)
- **Maximum depth of plant growth**
- **Community frequency of occurrence** (number of intercept points where aquatic plants were detected divided by the number of intercept points shallower than the maximum depth of plant growth)
- **Mean intercept point taxonomic richness** (the average number of taxa per intercept point)
- **Mean intercept point native taxonomic richness** (the average number of native taxa per intercept point)
- **Taxonomic frequency of occurrence within vegetated areas** (the number of intercept points where a particular taxon (e.g., genus, species, etc.) was detected divided by the total number of intercept points where vegetation was present)
- **Taxonomic frequency of occurrence at sites within the photic zone** (the number of intercept points where a particular taxon (e.g., genus, species, etc.) was detected divided by the total number of intercept points which are equal to or shallower than the maximum depth of plant growth)
- **Relative taxonomic frequency of occurrence** (the number of intercept points where a particular taxon (e.g., genus, species, etc.) was detected divided by the sum of all species' occurrences)
- **Mean density** (the sum of the density values for a particular species divided by the number of sampling sites)
- **Simpson Diversity Index (SDI)** is an indicator of aquatic plant community diversity. SDI is calculated by taking one minus the sum of the relative frequencies squared for each species present. Based upon the index of community diversity, the closer the SDI is to one, the greater the diversity within the population.
- **Floristic Quality Index (FQI)** (This method uses a predetermined [Coefficient of Conservatism](#) (C), that has been assigned to each native plant species in Wisconsin, based on that species' tolerance for disturbance. Non-native plants are not assigned conservatism coefficients. The aggregate conservatism of all the plants inhabiting a site determines its floristic quality. The mean C value for a given lake is the arithmetic mean of the coefficients of all native vascular plant species occurring on the entire site, without regard to dominance or frequency. The FQI value is the mean C times the square root of the total number of native species. This formula combines the conservatism of the species present with a measure of the species richness of the site.

4.3 SHORELINE CHARACTERIZATION

The point intercept method described above may not accurately identify emergent and floating leaved aquatic plants in near shore areas. Therefore, a boat tour was completed traveling the entire perimeter of the lake's shoreline. During the boat tour, visual observations of the emergent

and floating leaved plant communities were located and recorded. The boat tour also included a shoreline characterization, which provides an evaluation of shoreline development on the Lake. The following scale was used to rate the level of shoreline development.

- 1: Undeveloped** (i.e. Forested or wetland)
- 2: Minor development** (i.e. Properties may have mostly natural shoreline, sparse structures set further away from the lake, one pier, and little or no clearing of natural vegetation).
- 3: Moderate development** (i.e. Properties may exhibit clearing and/or manipulation to the shore and lawn areas but not to waters edge. More elaborate piers or boathouses may be present).
- 4: Major development** (i.e. Properties may include large lawn areas extending to the shoreline, which contains little or no natural shoreline vegetation. Increased building density, possibly close to the shore, multiple docks or boathouses, and significant shoreline alteration such as seawalls or rip rap may be present).

4.4 PUBLIC INVOLVEMENT, QUESTIONNAIRE, AND PLAN REVIEW

A public questionnaire was developed by Northern Environmental, the PITLC and the WDNR. This questionnaire was designed to gauge lake users' opinions on a number of important topics related to APM Plan implementation. The survey inquired about the users' perception of aquatic plant problems and other lake issues. The survey was also developed to determine what lake users consider an appropriate plant management intensity and cost. The public questionnaire was sent out to lake residents within the total PITLC project study area with their 2007 taxes in early 2008 and can be found in Appendix B.

Public involvement and education efforts included a presentation by Northern Environmental with the PITLC members on June 26, 2007 to discuss and kick off the APM Plan project. In November of 2007, draft copies of the APM Plan for Crab Lake, and other Presque Isle Town lakes, were submitted to the WDNR and the PITLC for distribution to, and comments from, lake residents. Comments from PITLC and the Vilas County Land and Water Conservation Department were received in December 2007. Edits to the documents were completed base on the comments received. Final preparation of the APM plan was on hold waiting WDNR review of the draft documents. Hearing no comments from the WDNR by April 8th, 2008, edited copies were bound and sent to the WDNR.

On May 3, 2008 a Northern Environmental project manger met with the PITLC, two members of the Vilas County Land and Water Department and the WDNR Lake Management Coordinator to discuss the findings of the APM Plans for the Presque Isle Town lakes. WDNR Lake Management Coordinator announced at that meeting that the WDNR review had not been completed and the group should anticipate comment yet to come. On December 22 of 2008, WDNR submitted 58 comments on the draft Crab APM plan. Northern Environmental Project Manager met with the PITLC, a representative of the Town of Presque Isle, Vilas County Land and Water Department Invasive Species Coordinator and the WDNR Lake Management Coordinator on January 21, 2009 to discuss the WDNR's comments. On April 17, 2009 Northern Environmental submitted a rewritten plan to the President of the PITLC for review and comment.

5.0 Discussion of Project Results

5.1 AQUATIC PLANT ECOLOGY

Aquatic plants are vital to the health of a water body. Unfortunately, people all too often refer to rooted aquatic plants as “weeds” and ultimately wish to eradicate them. This type of attitude, and the misconceptions it breeds, must be overcome in order to properly manage a lake ecosystem. Rooted aquatic plants (macrophytes) are extremely important for the well being of a lake community and possess many positive attributes. Despite their importance, aquatic macrophytes sometimes grow to nuisance levels that hamper recreational activities. This is especially prevalent in degraded ecosystems. The introduction of certain aquatic invasive species (AIS), such as EWM, often can exacerbate nuisance conditions, particularly when they compete successfully with native vegetation and occupy large portions of a lake.

When “managing” aquatic plants, it is important to maintain a well-balanced, stable, and diverse aquatic plant community that contains high percentages of desirable native species. To be effective, aquatic plant management in most lakes must maintain a plant community that is robust, species rich, and diverse. Appendix C1 includes a discussion about aquatic plant ecology, habitat types and relationships with water quality.

5.2 AQUATIC INVASIVE SPECIES

Aquatic Invasive Species (AIS) are aquatic plants and animals that have been introduced by human action to a location, area, or region where they did not previously exist. AIS often lack natural control mechanisms they may have had in their native ecosystem and may interfere with the native plant and animal interactions in their new “home”. Some AIS have aggressive reproductive potential and contribute to a decline of a lake’s ecology and interfere with recreational use of a lake. Common Wisconsin AIS include:

- Eurasian Watermilfoil
- Curly Leaf Pondweed
- Zebra Mussels
- Rusty Crayfish
- Spiny Water Flea
- Purple Loosestrife

Appendix C2 provides additional information on these AIS. Populations of rusty crayfish (*Orconectes rusticus*) have been identified by the WDNR within Crab Lake. Rusty crayfish are large crustaceans native to the Ohio River Valley that feed aggressively on aquatic plants, small invertebrates, small fish, and fish eggs.

5.3 2007 AQUATIC PLANT SURVEY

The survey was carried out August 2 – 4, 2007, and included sampling at 1204 intercept points. The aquatic macrophyte community of the Lake included 27 floating leaved, emergent, and submerged aquatic vascular plant species during 2007. Table 1 lists the taxa identified during the 2007 aquatic plant survey. Figures 4a through Figure 4g illustrate the locations of each species identified.

Vegetation was identified to a maximum depth of 23 feet (photic zone). Aquatic vegetation was detected at 15.4 percent (%) of photic zone intercept points. A diverse plant community inhabited the lake during 2007. The Simpson Diversity Index value of the community was 0.91. The taxonomic richness was 27 species and there was an average of 0.29 species identified at points that were within the photic zone. There was an average of 1.87 species present at points with vegetation present. Table 2 summarizes these overall aquatic plant community statistics.



Wild Celery
Source: UW Herbarium Website

The most abundant aquatic plant identified during the aquatic plant survey was wild celery (*Vallisneria americana*). It exhibited a 5.9% frequency of occurrence (percent of photic zone intercept points at which the taxa was detected). It was present at 38.2% of the sites with vegetation, and had a 20.4% relative frequency of occurrence. Table 3 includes the abundance statistics for each species.

Wild celery (*Vallisneria americana*) also known as eel-grass or tape-grass. It has ribbon-like leaves that tend to grow until they emerge in clusters along the waters surface. Wild celery is a premiere source of food for waterfowl. All portions of the plant are consumed. Beds of wild celery are also considered good fish habitat providing shade, shelter and feeding opportunities (Borman, et al., 1997).

Dwarf watermilfoil (*Myriophyllum tenellum*) was the second most abundant vascular plant species occurring at 3.6% of the photic zone. It was present at 23.5% of the sites with vegetation and had a 12.57% relative frequency of occurrence.

Dwarf watermilfoil (*Myriophyllum tenellum*) is very different in appearance from all other milfoils. It has short, unbranched stems on which the leaves are small scales or bumps along the stem. Dwarf water milfoil is found primarily on sandy substrate and in depths up to 12 feet. The plant provides good spawning habitat for panfish and the roots and rhizomes help to stabilize sediments (Borman, et al., 1997).



Dwarf watermilfoil
Source: UW Herbarium Website



Creeping spikerush
Source: UW Herbarium Website


Creeping spikerush (*Elocharis palustris*), large-leaf pondweed (*Potamogeton amplifoliosus*), and fern or Robbins pondweed (*Potamogeton robbinsii*) were the third most abundant vascular plant species occurring at 2.3% of the photic zone. The plants were present at 14.7% of the sites with vegetation and had a 7.9% relative frequency of occurrence.

Creeping spikerush (*Elocharis palustris*) has stems that rise either singly or in small clusters. Stems are topped with a spikelet that tapers to a point. Its shape resembles a burning match (Borman, et al., 1997). The spikelet is covered with scales. It prefers hard substrates (Nichols, 1999). The beds of this species provide food for a variety of waterfowl and helps anchor near shore sediments.

Large-leaf pondweed (*Potamogeton amplifolius*) is often referred to as musky weed or cabbage by anglers. Large leaf pondweed has robust stems and broad submersed leaves, which are slightly folded and lined with many veins. Floating leaves are oval on long stalks. It is found mainly in soft sediments in water one to several feet deep and is sensitive to increased turbidity. The plant is commonly grazed by waterfowl, offers habitat for invertebrates, and foraging opportunities for fish (Borman, e al., 1997).



Large-leaf Pondweed
Source: UW Herbarium Website



Fern Pondweed
Source: UW Herbarium Website

Fern or Robbins pondweed (*Potamogeton robbinsii*) is a submergent pondweed with robust stems and strongly two-ranked leaves, creating a feather or fern-like appearance while in the water. Fern pondweed sprouts in the spring and thrive in deeper water. Fern pondweed provides habitat for invertebrates that are grazed by waterfowl and also offers good cover for fish, particularly northern pike (Borman, et al., 1997).

5.3.1 FLOATING-LEAF PLANTS

The following three floating-leaf aquatic plant species were identified during the 2007 aquatic plant survey.

- *Brasenia schreberi* (watershield)
- *Nuphar variegata* (spatterdock)
- *Nymphaea odorata* (white water lily)

5.3.2 SUBMERGENT PLANTS

The following eighteen submergent aquatic plant species were identified during the 2007 aquatic plant survey.

- *Algae sp.* (filamentous algae) [algal]
- *Elodea canadensis* (elodea)
- *Eriocaulon aquaticum* (pipewort)
- *Fontinalis antipyretica* (common water moss)
- *Juncus pelocarpus* (brown-fruited rush)
- *Lobelia dortmana* (water lobelia)
- *Megalodonata beckii* (water marigold)
- *Myriophyllum alterniflorum* (alternate-flowered watermilfoil)
- *Myriophyllum tenellum* (dwarf watermilfoil)
- *Najas flexilis* (bushy pondweed or slender naiad)
- *Nitella sp.* (nitella) [algal]
- *Potamogeton amplifolius* (large-leaf pondweed)
- *Potamogeton epihydrus* (ribbon-leaf pondweed)
- *Potamogeton gramineus* (variable-leaf pondweed)
- *Potamogeton natans* (floating-leaf pondweed)

- *Potamogeton robbinsii* (fern or Robbins pondweed)
- *Potamogeton spirillus* (spiral-fruited pondweed)
- *Vallisneria americana* (wild celery)

5.3.3 EMERGENT PLANTS

The following six emergent aquatic plant species were identified during the 2007 aquatic plant survey.

- *Eleocharis palustris* (creeping spikerush)
- *Equisetum fluviatile* (water horsetail)
- *Pontederia cordata* (pickerelweed)
- *Schoenoplectus acutus* (hardstem bulrush)
- *Schoenoplectus subterminalis* (water bulrush)
- *Sparganium fluctuans* (floating-leaved bur-reed)

Table 1 lists the species identified. Appendix D includes brief descriptions of all aquatic plants identified.

5.4 NORTH CRAB LAKE PLANT SURVEY

A brief survey of North Crab Lake was conducted on August 5, 2008, with sampling at 14 points along the shoreline. Twenty-six aquatic plant species were identified including large purple bladderwort (*Utricularia purpurea*), a rare species in the State and currently listed as a species of concern.

5.4.1 FLOATING LEAF PLANTS

The following 3 floating-leaf aquatic plant species were identified during the 2007 aquatic plant survey.

- *Brasenia schreberi* (watershield)
- *Nuphar variegata* (spatterdock)
- *Nymphaea odorata* (white water lily)

5.4.2 SUBMERGENT PLANTS

The following 15 submergent aquatic plant species were identified during the 2007 survey.

- *Chara sp.* (muskgrass) [algal]
- *Elodea canadensis* (elodea)
- *Eriocaulon aquaticum* (pipewort)
- *Isoetes sp.* (quillwort)
- *Megalodonta beckii* (water marigold)
- *Myriophyllum alterniflorum* (alternate-flowered watermilfoil)
- *Myriophyllum tenellum* (dwarf watermilfoil)
- *Najas flexilis* (bushy pondweed or slender naiad)
- *Potamogeton amplifolius* (large-leaf pondweed)
- *Potamogeton epihydrus* (ribbon-leaf pondweed)
- *Potamogeton gramenius* (variable-leaf pondweed)
- *Potamogeton pusillus* (small pondweed)
- *Potamogeton robbinsii* (fern or Robbins pondweed)
- *Utricularia purpurea* (large purple bladderwort)
- *Vallisneria Americana* (wild celery)

5.4.3 EMERGENT PLANTS

The following 8 emergent aquatic plant species were identified during the 2007 aquatic plant survey.

- *Dulichium arundinaceum* (three-way sedge)
- *Eleocharis acicularis* (needle spikerush)
- *Eleocharis palustris* (creeping spikerush)
- *Equisetum fluviatile* (water horsetail)
- *Pontederia cordata* (pickerelweed)
- *Sagittaria sp.* (arrowhead)
- *Sparganium sp.* (bur-reed)
- *Typha latifolia* (broad-leaved cattail)

Table 5 lists the species identified. Appendix D includes brief descriptions of aquatic plants identified.

5.5 FLORISTIC QUALITY INDEX

Higher FQI numbers indicate higher floristic quality and biological integrity and a lower level of disturbance impacts. FQI varies around the state of Wisconsin and ranges from 3.0 to 44.6 with the average FQI of 22.2 (WDNR, 2005). The FQI calculated from the 2007 aquatic plant survey data was 37.07 and 34.09 for Crab and North Crab Lakes respectively. These FQI values are higher than the northern region average of 24.3 and suggest the Lakes exhibit very good water quality when using aquatic plants as an indicator. Tables 4 and 6 summarize the FQI values. Filamentous algae and common water moss do not have assigned values and were not included in calculation of the FQI. Arrowhead and bur-reed were not identified to species on North Crab Lake and were not included in that FQI calculation.

5.6 SHORELINE CHARACTERIZATION


Emergent and floating leaved plants identified along the shoreline outside of formal grid sample points included: *Sagittaria sp.* (arrowhead), *Nuphar variegata* (spatterdock), *Nymphaea odorata* (white water lily), *Eleocharis palustris* (creeping spikerush), *Pontederia cordata* (pickerelweed), *Typha sp.* (cattail), *Carex sp.* (sedges species) and *Sparganium sp.* (bur-reed). Refer to Appendix D for descriptions of these plants. Figure 5 illustrates the floating leaved and emergent plant locations identified during the boat survey. Plants identified during the shoreline survey but not during the point-intercept method were not included in the community statistics or calculation of the FQI.

The majority of the shoreline was moderately developed with residences. Some scattered blocks of undeveloped woodlands were present on the lakeshore. Figure 5 illustrates the level of shoreline development.

5.7 PUBLIC QUESTIONNAIRE (APPENDIX B)

There were 41 responses to the public questionnaire; 64% were Crab Lake shoreline landowners and 36% were nearby landowners on offshore property or another lake. Fishing, enjoying scenery, swimming, and canoeing/kayaking were the most popular activities enjoyed by survey respondents. Aquatic plant growth was reported to be a nuisance by the respondents at the following rate; 92% "rarely/never", 5% "sometimes", and 2% "most of the time/always". The major concerns of the respondents were water quality, fish populations and aquatic invasive species. Most of the respondents thought important goals of APM are to prevent the introduction of new AIS.

Respondents' APM Goals



The respondents reported having at least some awareness of the impacts of AIS to recreational use on a lake and were nearly split on their knowledge of AIS identification or methods for preventing the spread of AIS. These responses indicate that education will be a necessary component of APM.

Fifty-six percent of respondents believe that aquatic plant management is needed, 20% don't believe APM is needed and 24% are unsure. The majority of respondents (54%) thought that the level of appropriate aquatic plant management would be to manage problem areas. Most of the respondents thought that state and federal funding should be sought for AIS management activities and that user fees and room tax should be explored options.

5.8 WATER QUALITY SAMPLING

Citizen Lake-Monitoring collected water samples between 1993 and 2007 showed an average phosphorus reading of 11.4 ug/L and an average chlorophyll *a* reading of 2.54 ug/L. A Secchi reading were taken at the same time and averaged visible to 13.85 feet. All three results are concurrent with the oligotrophic status of the lake.

Along with water quality sampling, dissolved oxygen and temperature profiles were recorded at five foot intervals to a depth of 50 feet over the deepest portion of the lake. As lakes warm throughout the summer uneven heating of water occurs. This causes warmer water to "separate", or stratify, from cooler, denser water. This natural process is called stratification, which Crab Lake experiences. Between layers of stratified water a thermocline is established, which on Crab is at approximately 20 feet of depth, though it varies from year to year depending on the summer weather. During a hot summer, the thermocline will be found deeper than during an average, or cooler summer. A thermocline is a thin layer of water in which the temperature changes more rapidly with depth than in the layers above and below it. The area below the

thermocline may become low in oxygen concentration and limit the depth to which organisms can survive. At approximately 5.00 ppm of dissolved oxygen (DO), fish become stressed and at 2.00 ppm DO, most fish cannot survive. These levels occur at approximately 30 feet and 50 feet, respectively (Appendix I). Both before and after a thermocline develops, there is sufficient oxygen throughout the water column to support life.

6.0 Management Alternatives and Recommendations

Based on the goals of the stakeholders as mentioned in section 3.6, several management alternatives are available for this APM plan. Some general alternatives are discussed below. More information on management alternatives is included in Appendix E. Currently, the Northern Region of the WDNR is working under an aquatic plant management strategy that is officially titled Aquatic Plant Management Strategy, Northern Region WDNR, Summer, 2007 (working draft), or commonly referred to the NOR Region APM Strategy (Appendix G). This strategy lays out an approach for acceptable aquatic plant management in Northern Region lakes. The strategy protects native aquatic plant communities in northern Wisconsin and does not allow permits to control native plants unless documented circumstances of nuisance levels exist. The following management alternatives are based on the approaches described in the NOR Region APM Strategy, and incorporate recommendations of Bonestroo.

6.1 AQUATIC PLANT MAINTENANCE ALTERNATIVES

The maintenance alternative may be used at a lake in which a healthy aquatic plant community exists and invasive and non-native plant species are generally not present. The maintenance alternative is a protection-oriented management alternative because no significant plant problems exist or no active manipulation is required. This alternative can include an educational plan to inform lake shore owners of the value of a natural shoreline and encourage the protection of the lake water quality and the native aquatic plant community.

6.1.1 AQUATIC INVASIVE SPECIES MONITORING

No AIS were identified during the 2007 survey in Crab Lake. In order to monitor for AIS in the future a strong Citizen Lake Monitoring program that surveys for AIS is highly recommended. In some lake systems, native aquatic plants “hold their own” and AIS never grow to nuisance levels, in others, however, vigilant and active management is required. This can be based on several things including water quality. Data provided on the WDNR Citizen Lake Monitoring website indicates monitoring of water clarity is currently ongoing.

Assuming an AIS were to become established in the next several years, the most likely species would be EWM or CLP. If these or other AIS are found a sample should be collected and taken to the DNR for proper confirmation. The University of Wisconsin-Extension Lake’s Program provides training and coordinates the Citizen Lake Monitoring Program. More information about the program is available by contacting Laura Herman, Citizen Lake Monitoring Network Education Specialist, (715) 346-3989, email: lh Herman@uwsp.edu, website: <http://www.uwsp.edu/cnr/uwexlakes/clmn/>.

Bonestroo also recommends completing lake-wide aquatic plant surveys every 5 years (essentially repeating the 2007 point intercept aquatic plant survey) to monitor changes in the overall aquatic plant community and the effects of the APM activities. Aquatic plant communities may change with varying water levels, water clarity, nutrient levels, and aquatic plant management actions.

6.1.2 CLEAN BOATS/CLEAN WATERS CAMPAIGN

Measures for the prevention of the introduction of AIS to the lake should be a priority. To prevent the spread of AIS into Crab Lake, a monitoring program such as Clean Boats/Clean Waters is an excellent choice. There is one public landing on Crab Lake and lake residents are currently participating in Clean Boats/Clean Waters (CB/CW) program. This program is carried out by trained volunteers who inspect the incoming boats at public launches. Signage also accompanies the use of CB/CW to inform lake users of proper identification of AIS and boat inspection procedures. Education of the public, along with private property and resort owners, about inspecting watercraft for AIS before launching a boat or leaving access sites on other lakes could help prevent new AIS infestations. Contact with lake users at this time is a great way to distribute other educational materials. Continuation of this program is recommended and should be promoted by the current CB/CW coordinator on the lake.

6.1.3 AQUATIC PLANT PROTECTION AND SHORELINE MANAGEMENT

Protection of the native aquatic plant community is needed to slow the spread of EWM from lake to lake and within a lake once established. Therefore, riparian landowners should refrain from removing native vegetation. Additionally, EWM can thrive in nutrient (phosphorus and nitrogen) enriched waters or where nutrient rich sediments occur. Two simple actions can prevent excessive nutrients and sediments from reaching the lake.

The first activity is the restoration of natural shorelines, which act as a buffer for runoff containing nutrients and sediments. Properties classified in the shoreland survey as having a level 3: Moderate Development or level 4: Major Development, would be good candidates for shoreland restorations. Establishing natural shoreline vegetation can sometimes be as easy as not mowing to the waters edge. Native plants can also be purchased from nurseries for restoration efforts. Shoreline restoration has the added benefits of providing wildlife habitat and erosion prevention. A vegetated buffer area can also prevent surface water runoff from roads, parking areas and lawns from carrying nutrients to the lake.

The Vilas County Land and Water Conservation Department offers a cost-share program for county landowners. The primary emphasis of the program continues to be to restore native vegetation to shoreland property. For shoreline restoration projects and other conservation practices involving revegetation activities, landowners are reimbursed up to 70% of the costs of planting and purchasing native trees, shrubs, and wildflowers. Interested landowners can contact the Vilas County Land & Water Conservation Department at (715) 479-3648 to request an application form for the program. Another avenue to fund shoreland restoration is the WDNR Lake Protection Grant program. This program offers 75% of the project cost covered by the state up to \$200,000. For more information on the Lake Protection Grant program contact the Lake Management Coordinator at the WDNR Rhinelander Service Center by calling (715) 365-8937.

The second easy nutrient prevention effort is to use lawn fertilizers only when a soil test shows a lack of nutrients. Phosphorus free fertilizers should be used when possible. The fertilizers commonly used for lawns and gardens have three major plant macronutrients: Nitrogen, Phosphorus, and Potassium. These are summarized on the fertilizer package by three numbers. The middle number represents the amount of phosphorus. Since most Wisconsin lakes are "Phosphorus limited", meaning additions of phosphorus can cause increased aquatic plant or algae growth, preventing phosphorus from reaching the lake is a good practice. Landowners

should be encouraged to use phosphorus free fertilizers on lakeshore lawns. Local retailers and lawn care companies can provide soil test kits to determine a lawn's nutrient needs. Of course, properties with an intact natural buffer require very little maintenance, and no fertilizers.

Another possible source of nutrients to a lake is the septic systems surrounding the lake. Septic systems should be properly installed and maintained in order to prevent improperly treated wastewater, which carries a lot of nutrients, from reaching the lake. Property owners who are not sure if their septic system is adding nutrients to the lake should contact a professional inspector and have their system assessed.

6.1.4 PUBLIC EDUCATION AND INVOLVEMENT

The PITLC should continue to keep abreast of current AIS issues throughout the County. The County Land and Water Resource Conservation Department and the WDNR Lakes Coordinator, and the UW Extension are good sources of information. Many important materials can be ordered at the following website:

<http://www.uwsp.edu/cnr/uwexplakes/publications/>

Appendix G includes resources for further information about public education opportunities.

If the above hyperlink to web address becomes inactive, please contact Bonestroo for appropriate program and contact information.

6.2 AQUATIC PLANT MANIPULATION ALTERNATIVES

The management alternative may be used when aquatic plants present some sort of problem that must be dealt with or manipulated by human action. The WDNR NOR Region APM Strategy states "Newly-discovered infestations, if found on a lake with an approved lake management plan, the invasive species can be controlled via an amendment to the approved plan." The following alternatives are based on the assumption that the PITLC will meet in consultation with the WDNR before pursuing manipulation of AIS populations.

6.2.1 MANUAL REMOVAL

Native plants may be found at nuisance levels at individual properties. Manual removal efforts, including hand raking or hand pulling unwanted plants, are allowed under Wisconsin law, to a maximum width of 30 feet (recreational zone). The intent is to provide pier, boatlift or swimming raft access in the recreation zone. A permit is not required for hand pulling or raking if the maximum width cleared does not exceed this 30-foot recreation zone (manual removal of any native aquatic vegetation beyond the 30-foot area would require a permit from the WDNR that satisfies the requirements of Chapter NR 109, Wisconsin Administrative Code, see Appendix F). However manual removal is **not** recommended because it could open a niche for non-native invasive aquatic plants to occupy. Removal of native plants also destroys habitat for fish and wildlife.

If an Aquatic invasive plant is found in a small population hand pulling is a good first line of defense. If EWM or CLP ever becomes established within Crab Lake, manual (hand) removal of these plants in small, isolated populations, particularly in shallow water would be appropriate. No permit is required to remove non-native invasive aquatic vegetation, as long as the removal is

conducted completely by hand with no mechanical assistance of any kind. All aquatic plant material must be removed from the water to minimize dispersion and re-germination of unwanted aquatic plants. Portions of the roots may remain in the sediments, so removal may need to be repeated periodically throughout the growing season. Before significant plant removal is undertaken, a sample of the species assumed to be EWM or CLP should be brought to and confirmed by the WDNR.

Manual removal of aquatic plants can be quite labor intensive and time consuming. This technique is well suited for small areas in shallow water where property owners can weed the aquatic garden. Hiring laborers to remove aquatic vegetation is an option, but also increases cost. Scuba divers can be contracted to remove unwanted vegetation in deeper areas. Benefits of manual removal by property owners include low cost compared to chemical control methods, quick containment of pioneering (new) populations of invasive aquatic plants, and the ability for a property owner to slowly and consistently work on active management. The drawback of this alternative is that pulling aquatic plants include the challenge of working in the water, especially deep water, the threat of letting fragments escape and colonize a new area, and the fact that control of any significant sized population is quite labor intensive. Again, hiring laborers to remove aquatic vegetation is an option, but also increases cost.

Landowners removing plants manually should learn to identify the aquatic plant species. If an individual has questions about a particular aquatic plant or what level of manual removal is allowed, they should talk to the Vilas County Land & Water Conservation Department at (715) 479-3648, or the Kevin Gauthier, Lakes Management Coordinator, Wisconsin Department of Natural Resources, 107 Sutliff Ave, Rhinelander, (715) 365-8937. Appendix F includes additional resources for plant identification.

6.2.2 AQUATIC INVASIVE PLANT SPECIES CHEMICAL HERBICIDE TREATMENT

If Crab Lake becomes infested with EWM or CLP of areas of approximately ¼ acre or greater, a chemical herbicide treatment may be an appropriate way to conduct restoration of native plants. Before any specific course of action is undertaken the WDNR must be consulted. As of the time this report is written the consultation would begin with Kevin Gauthier, Lakes Management Coordinator in Rhinelander, (715) 365-8937. All herbicide treatments must be undertaken with a WDNR issued permit (NR 107 Wisconsin Administrative Code). A WDNR, AIS Early Detection and Rapid Response Grant is usually the best place for a lake group to receive financial assistance for chemical treatment of a newly discovered AIS population.

When using chemicals to control AIS it is a good idea to reevaluate the lake and the extent of the AIS conditions before, during and after chemical treatment. The WDNR may require another whole-lake plant survey and will certainly require a proposed treatment area survey. Along with the above mentioned survey, pre and post treatment monitoring should be included for all aquatic plant treatments and is typically a WDNR requirement in their Northern Region.

The science regarding what chemicals are most effective and how they can be used is constantly being updated. Currently EWM is the most common aquatic invasive plant species targeted for chemical treatment in the Northwoods. At present, granular 2,4-D is the most common herbicide used on EWM in the Northwood's area. In order to decrease damage to native plants and be as

selective as possible for EWM, treatments are completed in the spring when native plant growth is minimal.

Chemical treatment is usually a long term commitment and requires a specific plan with a goal set for "tolerable" levels of the relevant AIS. One such landmark might be 10% or less of the littoral area being occupied by aquatic invasive plants. WDNR recommends conducting a whole-lake point-intercept survey on a five year bases (for Crab Lake the next would be 2012). Such a survey may reveal a new AIS and at the very least would provide good trend data to see how the aquatic plant community is evolving.

Advantages of herbicides include broader control than hand pulling, and represents a true restoration effort, which harvesters do not (this is why harvesters are not discussed in this document). Disadvantages include negative public perception of chemicals in natural lakes, the potential to affect non-target plant species (if not applied at an appropriate application rate and/or time of year) and water use restrictions after application may be necessary.

7.0 Conclusion and Recommended Action Plan

No aquatic invasive plants were found during the aquatic plant survey in 2007. The fact that the native plant community had an above average plant FQI illustrates that there is a unique and diverse plant community in Crab Lake. Such a plant community is worthy of protection for human disturbance and the potential impact aquatic invasive species would have if introduced to this system. Because of that, the following recommended action plan focuses on maintenance of the current plant community and conservation of native plants.

7.1 RECOMMENDED ACTIVE GOALS

The recommended action plan includes actions for Crab Lake based on the Maintenance Alternative listed above in Section 6. The PITLC president has approved the following active goals. It will be up to residents of Crab Lake and the PITLC to determine the actions, find the funding, and gather the individuals needed to implement the active goals.

Active Goal: To continue and expand the WDNR Clean Boats, Clean Waters program on Crab Lake.

Active Goal: To provide visitors with educational information concerning the potential impact their activities could have on introductions aquatic invasive species, wildlife, habitats and Crab Lake water quality.

Active Goal: To implement and maintain an aquatic invasive species monitoring program that will survey for invasive species, and if any are found, monitor their locations and extent of populations spread.

Active Goal: To continue the Crab Lake comprehensive water quality monitoring program through the WDNR Citizen Lake Monitoring Network.

Active Goal: To work in concert with the WDNR staff and representatives of fishing related businesses to evaluate Crab Lake fish management practices and develop goals in order to maintain and enhance a quality family sport fishery.

Active Goal: To develop and implement a fishing tournament AIS management and monitoring program in order to be prepared if fishing tournaments come to Crab Lake.

Active Goal: To enforce the Town of Presque Isle's 200 foot no-wake areas (from shoreline and islands) ordinances in order to minimize recreational impacts on the plant community, shoreline habitats, and to promote safe boating.

Active Goal: To support the identification and preservation of critical species and critical habitat lands, and wetlands within the watershed. (These are areas with rare vegetation, important habitat for wildlife, or important spawning and nursery

areas for fish. Preservation of these lands has a direct impact on the water quality of the lake).

Active Goal: To provide education and information to shoreline property owners regarding how native aquatic plant protection and shoreline management can slow the spread of aquatic invasive plants (if they become introduced), improve the lake fishery, improve wildlife habitat and affect the quality of the water in the lake (including development of a *shoreline restoration packet* that could be given to landowners who's property has development categorized as Moderate or Major).

Active Goal: To encourage the incorporation of water quality protection measures in the design, construction and maintenance of all lake access sites on Crab Lake (e.g. storm water control, site drainage control, appropriate plant matter disposal, and watercraft wash down facilities if found to be needed). (*However, see Appendix J*)

Active Goal: To meet on a regular basis with local government agencies and representatives of lakes located within the Town of Presque Isle, to identify essential and new lake management issues and determine collaborative solutions.

7.2 CLOSING

This APM Plan was prepared in cooperation with the Presque Isle Town Lakes Committee. It includes the major components outlined in the WDNR Aquatic Plant Management guidance. The "Recommended Action Plan" section of this report can be used as a stand alone document to facilitate EWM management activities for the lake. This section outlines important monitoring and management activities. The greater APM Plan document and appendices provides a central source of information for the lake's aquatic plant community information, the overall lake ecology, and sources of additional information. If there are any questions about how to use this APM Plan or its contents, please contact Bonestroo.

This APM Plan should be updated periodically to reflect current aquatic plant problems, and the most recent acceptable APM methods. Information regarding aquatic plant management and protection is available from the WDNR website:

<http://dnr.wi.gov/org/water/fhp/lakes/aquaplan.htm> or from Bonestroo upon request.

8.0 References

While not all references are specifically cited, the following resources were used in preparation of this report.

Borman, Susan, Robert Korth, and Jo Temte, *Through the Looking Glass, A Field Guide to Aquatic Plants*, Wisconsin Lakes Partnership, 1997

Carlson, R. E., A trophic state index for lakes. *Limnology and Oceanography*, 22:361-369, 1977

Fassett, Norman C., *A Manual of Aquatic Plants*, The University of Wisconsin Press, Madison, Wisconsin, 1975

Getsinger, Kurt D., and H.E. Westerdahl, *Aquatic Plant Identification and Herbicide Use Guide, Volume II Aquatic Plants and Susceptibility to Herbicides*, U.S. Northern Environmental Waterways Experiments Station, Technical Report A-88-9, 1988

Jester, Laura, Bozek, Michael, Helsel, Daniel, and Sheldon, Sallie, *Euhrychiopsis lecontei Distribution, Abundance, and Experimental Augmentation for Eurasian watermilfoil Control in Wisconsin Lakes*, *Journal Aquatic Plant Management*, 38:88-97

Madsen, John, *Point Intercept and Line Intercept Methods for Aquatic Plant Management, Aquatic Plant Control Technical Note MI-02*, February 1999

Nichols, Stanley A. *Distribution and habitat descriptions of Wisconsin lake plants*, Wisconsin Geological and Natural History Survey Bulletin 96, 1999

North America Lake Management Society of Aquatic Plant Management Society (NALMS), *Aquatic Plant Management in Lakes and Reservoirs*, 1997

Prescott, G.W., *How to Know the Aquatic Plants*, Wm. C. Brown Publishers, Dubuque, Iowa, 1980

United States Department of Agriculture, *Soil Survey of Vilas County, Wisconsin*. 1988

United States Geological Survey, *Heafford Junction, Wisconsin Quadrangle, 7.5 minute (Topographic) Series*, 1982

United States Geological Survey, Nonindigenous Aquatic Species, (<http://nas.er.usgs.gov/queries/collectioninfo.asp?>), Accessed November 13, 2007

Welsh, Jeff, *Guide to Wisconsin Aquatic Plants*, Wisconsin Department of Natural Resources Publication WR 173, 1992 revised

Wetzel, Robert G., *Limnology*, 1983

Wisconsin Department of Natural Resources, *Aquatic Plant Management in Wisconsin DRAFT*, April 25 2005

Wisconsin Department of Natural Resources, *Aquatic Invasive Species Website* (<http://dnr.wi.gov/invasives/aquatic/>), Accessed April 2007

Wisconsin Department of Natural Resources, *Listing of Wisconsin Waters with Eurasian Water-Milfoil (current as of 01/02/2007)*, 2007

Wisconsin Department of Natural Resources, *Fish Stocking Website*

(http://infotrek.er.usgs.gov/doc/wdnr_biology/Public_Stocking/StateMapHotspotsAllYears.htm), Accessed April, 2007

Wisconsin Department of Natural Resources, *Wisconsin Lakes*, Publication # PUB-FH-800, 2005

Figures

Tables

Appendix A - Point Intercept Sample Coordinates

Appendix B – Summary of Public Survey

Appendix C1 – Importance of Aquatic Plants to Lake Ecosystem

AQUATIC PLANT TYPES AND HABITAT

Aquatic plants can be divided into two major groups: microphytes (phytoplankton and epiphytes) composed mostly of single-celled algae, and macrophytes that include macro algae, flowering vascular plants, and aquatic mosses and ferns. Wide varieties of microphytes co-inhabit all habitable areas of a lake. Their abundance depends on light, nutrient availability, and other ecological factors.

In contrast, macrophytes are predominantly found in distinct habitats located in the littoral (i.e., shallow near shore) zone where light sufficient for photosynthesis can penetrate to the lake bottom. The littoral zone is subdivided into four distinct transitional zones: the eulittoral, upper littoral, middle littoral, and lower littoral (Wetzel, 1983).


Eulittoral Zone: Includes the area between the highest and lowest seasonal water levels, and often contains many wetland plants.

Upper Littoral Zone: Dominated by emergent macrophytes and extends from the shoreline edge to water depths between 3 and 6 feet.

Middle Littoral Zone: Occupies water depths of 3 to 9 feet, extending deeper from the upper littoral zone. The middle littoral zone is often dominated by floating-leaf plants.

Lower Littoral Zone: Extends to a depth equivalent to the limit of the photic zone, which is the maximum depth that sufficient light can support photosynthesis. This area is dominated by submergent aquatic plant types.

The following illustration depicts these particular zones and aquatic plant communities.



Aquatic Plant Communities Schematic



The abundance and distribution of aquatic macrophytes are controlled by light availability, lake trophic status as it relates to nutrients and water chemistry, sediment characteristics, and wind energy. Lake morphology and watershed characteristics relate to these factors independently and in combination (NALMS, 1997).

AQUATIC PLANTS AND WATER QUALITY

In many instances aquatic plants serve as indicators of water quality due to the sensitive nature of plants to water quality parameters such as water clarity and nutrient levels. To grow, aquatic plants must have adequate supplies of nutrients. Microphytes and free-floating macrophytes (e.g., duckweed) derive all their nutrients directly from the water. Rooted macrophytes can absorb nutrients from water and/or sediment. Therefore, the growth of phytoplankton and free-floating aquatic plants is regulated by the supply of critical available nutrients in the water column. In contrast, rooted aquatic plants can normally continue to grow in nutrient-poor water if lake sediment contains adequate nutrient concentrations. Nutrients removed by rooted macrophytes from the lake bottom may be returned to the water column when the plants die. Consequently, killing too many aquatic macrophytes may increase nutrients available for algal growth.

In general, an inverse relationship exists between water clarity and macrophyte growth. That is, water clarity is usually improved with increasing abundance of aquatic macrophytes. Two possible explanations are postulated. The first is that the macrophytes and epiphytes out-compete phytoplankton for available nutrients. Epiphytes derive essentially all of their nutrient needs from the water column. The other explanation is that aquatic macrophytes stabilize bottom sediment and limit water circulation, preventing re-suspension of solids and nutrients (NALMS, 1997).

If aquatic macrophyte abundance is reduced, then water clarity may suffer. Water clarity reductions can further reduce the vigor of macrophytes by restricting light penetration. Studies have shown that if 30 percent or less of a lake areas occupied by aquatic plants is controlled, water clarity will generally not be affected. However, lake water clarity will likely be reduced if 50 percent or more of the macrophytes are controlled (NALMS, 1997).

Aquatic plants also play a key role in the ecology of a lake system. Aquatic plants provide food and shelter for fish, wildlife and invertebrates. Plants also improve water quality by protecting shorelines and the lake bottom, improving water quality, adding to the aesthetic quality of the lake and impacting recreational activities.

Appendix C2 – Aquatic Invasive Species

INVASIVE AQUATIC PLANTS

Invasive species have invaded our backyards, forests, prairies, wetlands, and waters. Invasive species are often transplanted from other regions, even from across the globe. "A species is regarded as invasive if it has been introduced by human action to a location, area, or region where it did not previously occur naturally (i.e., is not native), becomes capable of establishing a breeding population in the new location without further intervention by humans, and spreads widely throughout the new location " (Source: WDNR website, Invasive Species, 2007). AIS include plants and animals that affect our lakes, rivers, and wetlands in negative ways. Once in their new environment, AIS often lack natural control mechanisms they may have had in their native ecosystem and may interfere with the native plant and animal interactions in their new "home". Some AIS have aggressive reproductive potential and contribute to ecological declines and problems for water based recreation and local economies. AIS often quickly become a problem in already disturbed lake ecosystems (i.e. one with relatively few native plant species). While native plants provide numerous benefits, AIS can contribute to ecological decline and financial constraints to manage problem infestations.

Eurasian Watermilfoil (*Myriophyllum spicatum*)

EWM is the most common AIS found in Wisconsin lakes. EWM was first discovered in southeast Wisconsin in the 1960's. During the 1980's, EWM began to spread to other lakes in southern Wisconsin and by 1993 it was common in 39 Wisconsin counties. EWM continues to spread across Wisconsin and is now found in the far northern portion of the state including Vilas County.

Unlike many other plants, EWM does not rely on seed for reproduction. Its seeds germinate poorly under natural conditions. It reproduces vegetatively by fragmentation, allowing it to disperse over long distances. The plant produces fragments after fruiting once or twice during the summer. These shoots may then be carried downstream by water currents or inadvertently picked up by boaters. EWM is readily dispersed by boats, motors, trailers, bilges, live wells, or bait buckets, and can stay alive for weeks if kept moist (WDNR website, 2007).

Once established in an aquatic community, EWM reproduces from shoot fragments and stolons (runners that creep along the lake bed). As an opportunistic species, EWM is adapted for rapid growth early in spring. Stolons, lower stems, and roots persist over winter and store the carbohydrates that help milfoil claim the water column early in spring, photosynthesize, divide, and form a dense leaf canopy that shades out native aquatic plants. Its ability to spread rapidly by fragmentation and effectively block out sunlight needed for native plant growth often results in monotypic stands. Monotypic stands of EWM provide only a single habitat, and threaten the integrity of aquatic communities in a number of ways; for example, dense stands disrupt predator-prey relationships by fencing out larger fish, and reducing the number of nutrient-rich native plants available for waterfowl (WDNR website, 2007).



Dense stands of EWM also inhibit recreational uses like swimming, boating, and fishing. The visual impact that greets the lake user on milfoil-dominated lakes is the flat yellow-green of matted vegetation, often prompting the perception that the lake is "infested" or "dead". Cycling of nutrients from sediments to the water column by EWM may lead to deteriorating water quality and algae blooms of infested lakes (WDNR website, 2007).

Curly leaf pondweed (*Potamogeton crispus*)

Curly-leaf pondweed (CLP) spreads through burr-like winter buds (turions), which are moved among waterways. These plants can also reproduce by seed, but this plays a relatively small role compared to the vegetative reproduction through turions. New plants form under the ice in winter, making CLP one of the first nuisance aquatic plants to emerge in the spring.

The leaves of curly-leaf pondweed are reddish-green, oblong, and about 3 inches long, with distinct wavy edges that are finely toothed. The stem of the plant is flat, reddish-brown and grows from 1 to 3 feet long. The plant usually drops to the lake bottom by early July.



CLP becomes invasive in some areas because of its tolerance for low light and low water temperatures. These tolerances allow it to get a head start on and out-compete native plants in the spring. CLP forms surface mats that interfere with aquatic recreation in mid-summer, when most aquatic plants are growing, CLP plants are dying off. Plant die-offs may result in a critical loss of dissolved oxygen. Furthermore, the decaying plants can increase nutrients which contribute to algal blooms, as well as create unpleasant stinking messes on beaches (WDNR website, 2007).



Purple Loosestrife (*Lythrum salicaria*)

Purple loosestrife is a perennial herb 3-7 feet tall with a dense bushy growth form. Showy flowers vary from purple to magenta, possess 5-6 petals aggregated into numerous long spikes, and bloom from July to September. Leaves are opposite, nearly linear, and attached to four-sided stems without stalks. It has a large, woody taproot with fibrous rhizomes that form a dense mat.

Purple loosestrife was first detected in Wisconsin in the early 1930's, but remained uncommon until the 1970's. It is now widely dispersed in the state, and has been recorded in 70 of Wisconsin's 72 counties. Low densities in most areas of the state suggest that the plant is still in the pioneering stage of establishment. Areas of heaviest infestation are sections of the Wisconsin River, the extreme southeastern part of the state, and the Wolf and Fox River drainage systems.



This plant's optimal habitat includes marshes, stream margins, alluvial flood plains, sedge meadows, and wet prairies. It is tolerant of moist soil and shallow water sites such as pastures and meadows, although established plants can tolerate drier conditions. Purple loosestrife has also been planted in lawns and gardens, which is often how it has been introduced to many of our wetlands, lakes, and rivers. Purple loosestrife spreads mainly by seed, but it can also spread vegetatively from root or stem segments. A single stalk can produce from 100,000 to 300,000

seeds per year. Seed survival is up to 60-70%, resulting in an extensive seed bank. Mature plants with up to 50 shoots grow over 2 meters high and produce more than two million seeds a year. Germination is restricted to open, wet soils and requires high temperatures, but seeds remain viable in the soil for many years. Even seeds submerged in water can live for approximately 20 months (WDNR website, 2007).

OTHER AQUATIC INVASIVE SPECIES

The following AIS are not plants, but are mentioned here because they also can significantly disrupt healthy aquatic ecosystems.

Rusty Crayfish (*Orconectes rusticus*) are large crustaceans that feed aggressively on aquatic plants, small invertebrates, small fish, and fish eggs. They can remove nearly all the aquatic vegetation from a lake, offsetting the balance of a lake ecosystem. More information about this invader can be found at <http://dnr.wi.gov/invasives/fact/rusty.htm>.

Zebra Mussels (*Dreissena polymorpha*) are small freshwater clams that can attach to hard substrates in water bodies, often forming large of thousands of individual mussels. They are prolific filter feeders, removing valuable phytoplankton from the water, which is the base of the food chain in an aquatic ecosystem. More information about this invader can be found at

<http://dnr.wi.gov/invasives/fact/zebra.htm>.

Spiny Water Fleas (*Bythotrephes cederstoemi*) are predatory zooplankton (tiny aquatic animals) that have a barbed tail making up most of their body length (one centimeter average). They compete with small fish for food supplies (zooplankton) and small fish cannot swallow the spiny water flea due to the long spiny appendage. More research is being completed to determine the potential impacts of the spiny water flea. More information about this invader can be found at

<http://dnr.wi.gov/invasives/fact/spiny.htm>.

Appendix D – Descriptions of Aquatic Plants

Appendix E – Summary of Aquatic Plant Management Alternatives

Appendix F – NR 107 and NR 109 Wisconsin Administrative Code

Appendix G – Resource for Additional Information

Appendix H – Aquatic Plant Management Strategy, Northern Region WDNR, Summer, 2007 (working draft)

Appendix I – Temperature and Dissolved Oxygen Profiles

Appendix J – Additional Information / Lake Resident Comments

ADDITIONAL INFORMATION / LAKE RESIDENT COMMENTS

Crab Lake residents are interested and committed in preservation and protection. However, recent WDNR proposed logging actions will be at odds to their stated interest in preservation of the watershed.

- Close to the south shore, between the South Crab Lake Road and to within 100 feet of the lake the WDNR plans to let a contract for clear cut logging in November 2010. This will clear cut everything on state owned land, including all ephemeral ponds, swamps, and lands with steep vertical topography. Of special concern is the crown of trees on and either side of the sub-continental divide.
- The Crab Lake Property Owners Association has protested, proposed some modification to the original logging plan, and brought the matter to the attention of both the DNR citizen's oversight board and to Matt Frank, the secretary of the DNR.
- The DNR has, to date, not entertained any negotiated settlement proposed by the Lake Association to preserve the trees in any of the wetland areas nor on the sub-continental divide.
- Crab Lake residents strongly believe that these DNR proposed logging actions will be at opposition to their stated interest in preservation of the watershed.

Current Update - The DNR let the contract in fall 2010 for logging to be performed over the winter of 2011/12 or 2012/13, the time at the discretion of the logger. Our Ass'n continues to press the DNR in that their logging plan is in conflict with both the water management people (Kevin Gauthier's staff re: watershed preservation) and with their own 'Tree Retention and Managed Forest Law (August 3, 2011)' which talks about tree retention and correct silviculture practices as noted in the DNR Silviculture Handbook. Our Association is trying for another review/revision/appeal to the amount of logging but since the contract has been bid and let, the trees marked, we don't hold much hope.