

# Using dissolved carbon dioxide (CO<sub>2</sub>) to alter the behavior of round goby

## Background

This project aimed to determine whether elevated CO<sub>2</sub> levels in water could be effectively used as a behavioral deterrent to reduce the spread of invasive round goby. Elevated CO<sub>2</sub> concentrations in water may cause avoidance behavior and loss of equilibrium in fish. The research goal of this study was to identify the CO<sub>2</sub> concentrations that resulted in round goby avoidance behavior and loss of equilibrium as a first step to assess its potential effectiveness as a deterrent/barrier.



## Round Goby

Round goby (*Neogobius melanostomus*) is a non-native bottom-dwelling fish that was first detected in the Great Lakes in the 1990s. More recently, round goby were verified directly downstream of the Lake Winnebago system in Little Lake Butte des Morts. These aggressive fish can negatively affect native fish populations and disrupt the food chain. Round goby are listed as restricted under ch. NR 40.

Photo: Peter Van der Sluijs, license CC BY-SA 3.0

## Research Methods

- Scientists from the U.S. Geological Survey and the University of Illinois conducted laboratory experiments to assess how round goby responded to different levels of dissolved CO<sub>2</sub> in the water.
- Round goby were placed in one of two connected tanks. CO<sub>2</sub> was then added to the tank containing the fish to observe fish behavior and determine CO<sub>2</sub> concentrations when fish moved into the adjoining refuge tank (where no CO<sub>2</sub> was added).
- In separate trials, CO<sub>2</sub> was continuously added into a tank until fish were physically immobilized to determine concentrations that resulted in loss of equilibrium.



Peter van der Sluijs

## Summary of Results

- Round goby exhibited similar behavior responses to CO<sub>2</sub> as were seen in previous studies conducted on non-native Asian carp.
- Round goby voluntarily avoided high-CO<sub>2</sub> waters at a concentration threshold of 99 – 169 mg/L CO<sub>2</sub>.
- Round goby involuntarily lost equilibrium at 197 – 280 mg/L CO<sub>2</sub>.
- Results indicate that temperature plays a role in the effectiveness of this method. Dissolved CO<sub>2</sub> may be a more effective deterrent at low water temperatures.

## Potential applications:

Fish voluntarily avoided CO<sub>2</sub>-enriched water before it caused them to lose equilibrium and become unable to right themselves. These results suggest that with further study, it may be possible to use CO<sub>2</sub> as a deterrent for this species in place of a physical barrier. Additional research is needed on potential impacts of increased CO<sub>2</sub> on other non-target species. However, these laboratory experiments can inform future field trials on the use of CO<sub>2</sub> as a method to deter the continued spread of round goby and protect native fisheries and ecosystems.

## Related Publications and Resources

Cupp, A.R. et al. 2017. Using dissolved carbon dioxide to alter the behavior of invasive round goby. *Management of Biological Invasions*. 8(4): 567 – 574. [doi.org/10.3391/mbi.2017.8.4.12](https://doi.org/10.3391/mbi.2017.8.4.12)

This project was supported through donations to aquatic invasive species research. To learn more or donate, visit [www.dnr.wi.gov/lakes/sayyestolakes](http://www.dnr.wi.gov/lakes/sayyestolakes)

